(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 12 April 2001 (12.04.2001)

PCT

(10) International Publication Number WO 01/24781 A2

(51) International Patent Classification⁷: A61K 31/00

(21) International Application Number: PCT/CA00/01188

(22) International Filing Date: 6 October 2000 (06.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

2,285,690 7 October 1999 (07.10.1999) CA 60/158,043 7 October 1999 (07.10.1999) US 60/217,765 12 July 2000 (12.07.2000) US

- (71) Applicant (for all designated States except US): NOVA-NEURON INC. [CA/CA]; 5859 University Avenue, Sir Charles Tupper Medical Bldg., Room 15D7, Halifax, Nova Scotia B3H 4H7 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ROBERTSON, Harold, A. [CA/CA]; 2384 Clifton Street, Halifax, Nova Scotia B3K 4V1 (CA). DENOVAN-WRIGHT, Eileen, M. [CA/CA]; 22 Fleming Drive, Halifax, Nova Scotia B3P 1A9 (CA).

- (74) Agent: HELLER, David, J.; Ridout & Maybee, One Queen Street East, Suite 2400, Toronto, Ontario M5C 3B1 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

 Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GENE NECESSARY FOR STRIATAL FUNCTION, USES THEREOF, AND COMPOUNDS FOR MODULATING SAME

(57) Abstract: PDE10A, a gene that is normally highly expressed in mammalian striatum and elsewhere, has been found to decrease in expression during the development of CAG repeat disorders such as Huntington's disease. The invention teaches a method for detecting the presence of or the predisposition for a CAG repeat disorder. Compounds which modulate CAG repeat disorders and their uses are taught. Methods for screening for further compounds to modulate CAG repeat disorders are also taught.

Gene Necessary for Striatal Function, Uses Thereof, and Compounds for Modulating Same

CROSS-REFERENCE

This patent claims priority from Canadian Patent application no. 2,285,690 filed October 7, 1999, US provisional application no. 60/158,043 filed October 7, 1999, and US provisional application no. 60/217,765 filed July 12, 2000, entitled Gene Necessary for Striatal Function, Uses Thereof, and Compounds for Modulating Same.

FIELD OF THE INVENTION

The present invention relates to a polynucleotide, PDE10A, which is down-regulated during the development of CAG repeat disorders, such as Huntington's disease. The present invention also describes compounds that modulate CAG repeat disorders, processes for expressing PDE10A, and its agonists and antagonists, and uses of PDE10A, and its variants, derivatives, agonists and antagonists.

BACKGROUND OF THE INVENTION

Very few if any effective treatments exist for neurological disorders characterized by progressive cell loss, known as neurodegenerative diseases, as well as those involving acute cell loss, such as stroke and trauma.

Huntington's disease (HD) is an inherited neurological disorder that is transmitted in

autosomal dominant fashion. HD results from genetically programmed degeneration of neurons in certain areas of the brain. Huntington's disease is caused by a mutation of the gene *IT-15* that codes for the protein huntingtin. The huntingtin gene contains a polymorphic stretch of repeated CAG trinucleotides that encode a polyglutamine tract within huntingtin. If this tract exceeds 35 in number, Huntington's disease results. Huntington's disease is only one of a number of neurological diseases which are characterised by these polyglutamine repeats (Ross, 1997). Schizophrenia, Alzheimer's disease, stroke, trauma, and Parkinson's disease also affect the basal ganglia.

Huntingtin has no sequence similarity to known proteins (Group THDCR, 1993; Sisodia, 1998). The function of the normal or mutated HD form of huntingtin has not been defined by the prior art. It is evident, however, that the expression of the HD form of huntingtin leads to progressive and selective neuronal loss. It has been demonstrated that the GABA- and enkephalin-containing medium spiny projection neurons of the caudate-putamen eventually die as a result of HD (Richfield et al., 1994). Patients with minimal cell loss, however, still present with motor and cognitive symptoms suggesting that neuronal dysfunction, and not simply cell loss, contribute to the symptoms of HD. The motor symptoms of HD include the development of chorea, dystonia, bradykinesia and tremors (Young et al., 1986). Voluntary movements may also be affected such that there may be disturbances in speech (Ludlow et al., 1987) and degradation of fine motor co-ordination (Young et al., 1986). In addition to motor decline, emotional disturbances and cognitive loss are also evident during the progression of HD (Caine et al., 1978).

Despite the fact that huntingtin is ubiquitously expressed, HD specifically affects cells of the

basal ganglia, structures deep within the brain that have a number of important functions, including co-ordinating movement. The basal ganglia includes the caudate nucleus, the putamen, the nucleus accumbens and the olfactory tubercule. HD also affects the brain's outer surface, or cortex, which controls thought, perception, and memory. The mechanism by which only a small group of neurons in the striatum and cortex are rendered vulnerable to this ubiquitously expressed mutant protein is not known. There are no effective treatments for Huntington's disease.

Huntington's disease is widely believed to be a gain-of function disorder but neither the normal function nor the gained function of huntingtin is known. Because the function for huntingtin is not known, there is little insight into the disease process. It was believed that huntingtin was related to neuronal intranuclear inclusions (NII). However, recent results have cast doubt on our understanding of the role of the NII in Huntington's disease (Saudou et al., 1998) or in other CAG repeat disorders (Klement et al., 1998; see also commentary by Sisodia, 1998).

The development of a mouse carrying the 5' end of the human Huntington's disease gene (the promoter and first exon; Mangiarini et al., 1996) was an important step in the development of the tools that will allow us to understand the function (and gain-of-function) associated with huntingtin. R6/2 mice exhibit a rapidly progressing neurological phenotype with onset at about 8 weeks. This phenotype includes a movement disorder characterised by shuddering, resting tremor, epileptic seizures and stereotyped behaviour. These symptoms suggest that the function of the basal ganglia is affected by the expression of the human exon 1 transgene prior to neuronal cell death. By 12 weeks the affected mice have significantly reduced brain

weights and they die by about 13 weeks of age. Neuronal intranuclear inclusions (NII) develop at about 4 weeks (Davies et al., 1997). As is observed in human Huntington's disease patient, the R6/2 mice show changes in neuronal receptors (Cha et al., 1998). The present inventors have also demonstrated that changes in the expression of DARPP-32 and cannabinoid receptors change over time in HD mice; such changes have also been observed in human Huntington's disease patients (unpublished results). The loss of the cannabinoid receptor is one of the earliest documented changes that occur prior to neuronal degeneration in human HD patients. The R6/2 model, therefore, mimics the early phases of HD; a point in disease development where intervention would be most appropriate.

Human PDE10 was recently identified by identification of cDNA fragments published on the National Center for Biotechnology Information (NCBI) Expressed Sequence Tags (EST) database (Loughney et al., WO99/42596). While PDE10 was found to share homology with known PDEs, no function could be identified for PDE10.

SUMMARY OF THE INVENTION

The present invention provides the function and uses of a nucleotide segment, PDE10A, and compounds which inhibit or promote the development of CAG repeat disorders such as Huntington's Disease.

The invention teaches a method for identifying a compound which inhibits or promotes a CAG repeat disorder, comprising the steps of: (a) selecting a control animal having PDE10A and a test animal having PDE10A; (b) treating said test animal using a compound; and (c)

determining the relative quantity of RNA corresponding to PDE10A, as between said animals. In an embodiment, the animal is a mammal, preferably a mouse, and preferably a transgenic mouse. In an embodiment, the CAG repeat disorder is Huntington's disease.

The invention also teaches a method for identifying a compound which inhibits or promotes a CAG repeat disorder, comprising the steps of: (a) selecting a host cell containing PDE10A; (b) cloning said host cell and separating said clones into a test group and a control group; (c) treating said test group using a compound; and (c) determining the relative quantity of RNA corresponding to PDE10A, as between said test group and said control group. In an embodiment, the CAG repeat disorder is Huntington's disease.

The invention further teaches a method for detecting the presence of or the predisposition for a CAG repeat disorder, said method comprising determining the level of expression of RNA corresponding to PDE10A in an individual relative to a predetermined control level of expression, wherein a decreased expression of said RNA as compared to said control is indicative of a CAG repeat disorder. Preferably, the expression is measured by in situ hybridization, fluorescent in situ hybridization, polymerase chain reaction, or DNA fingerprinting technique. In an embodiment, the CAG repeat disorder is Huntington's disease.

The invention further teaches compositions for treating a CAG repeat disorder comprising a compound which modulates PDE10 expression and a pharmaceutically acceptable carrier.

The compound can be selected from the group consisting of: quinpirole, alloxan, miconazole nitrate, MDL-12330A and tetracyline derivatives such as demeclocycline. The compound

may be selected from the group consisting of: (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-methyl-pyrazino[2', 1':6,1]pyrido[3,4-b]indole-1,4-dione,

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-pyrazino[2',1':6,1]pyrido[3,4-]indole-1,4-dione, (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-isopropyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione, (3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-3-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione, and

(3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2,3-dimethyl-pyraz

ino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione, or from the group consisting of: KS-505,

IC224,SCH 51866, IBMX and Dipyridamole. The disorder can be HD.

The invention also teaches the use of a composition which modulates PDE10 for treating a CAG repeat disorder comprising administering the composition to a subject in need of such treatment, and such use of the composition which modulates PDE10 for treating HD.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a portion of an autoradiogram of the differential display reaction identifying PDE10A in mouse brain mRNA.

FIG. 2 is a northern blot confirming that PDE10A has a lower steady-state level of expression in the striatum of transgenic HD mice.

FIG. 3 is a nucleotide sequence of the differential display cDNA fragment of pPDE10A.

FIG. 4 shows the *in situ* hybridization of probe 1 to coronal and saggital brain sections of 10 week-old wild-type and HD mice.

FIG. 5 shows the *in situ* hybridization corresponding to spatial and temporal expression of PDE10A in brain sections of wild-type and HD mice over the period of time that the HD mice develop abnormal movements and postures.

FIG. 6 shows the *in situ* hybridization corresponding to expression of PDE10A in brain sections of one day old wild-type and HD mice.

FIG. 7 shows the *in situ* hybridization corresponding to distribution of the mRNA of PDE10A in mouse striatal neurons.

FIG. 8 is the *in situ* hybridization corresponding to mRNA distribution of the rat homologue of PDE10A in rat brain tissue.

FIG. 9 shows a Southern blot analysis of DNA from wild-type and transgenic HD mice hybridized to the pPDE10A cDNA probe.

FIG. 10 is a nucleotide sequence of cPDE10-1, and corresponds to SEQ ID NO. 1.

FIG. 11 is a restriction map of cPDE10-1.

FIG. 12 is a nucleotide sequence of cPDE10-2, and corresponds to SEO ID NO. 2.

FIG. 13 is a restriction map of cPDE10-2.

FIG. 14 is a schematic diagram showing the alignment of cPDE10-1 and -2 and the regions that are identical and unique between the two clones.

FIG. 15 is a nucleotide sequence of cPDE10A and RACEs, corresponding to SEQ ID NO. 11.

FIG. 16 is a map of PDE10A coding sequence and restriction sites.

FIG. 17 is a map of PDE10A coding sequence and features.

FIG. 18 is a restriction map of PDE10A.

FIG. 19 is a nucleotide sequence of cPDE10A and corresponds to SEQ ID NO. 12.

DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

The following illustrative explanations are provided to facilitate understanding of certain terms used frequently herein. The explanations are provided as a convenience and are not limitative of the invention.

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity", "similarity" or "homologous", as used in the art, are relationships between two or

more polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. Both identity and similarity can be readily calculated (Lesk, A. M., 1988; Smith, D. W., 1993; Griffin, A. M., and Griffin, H. G., 1994; von Heinje, G., 1987; and Gribskov, M. and Devereux, J., 1991). While there exist a number of methods to measure identity and similarity between two polynucleotide sequences, both terms are well known to skilled artisans (von Heinje, G., 1987; Gribskov, M. and Devereux, 1991; and Carillo, H., and Lipman, D., 1988). Methods commonly employed to determine identity or similarity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D. (1988). Methods to determine identity and similarity are codified in computer programs. Computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., 1984), BLASTP, BLASTN, and FASTA (Atschul, S. F. et al., 1990).

"Isolated" means altered "by the hand of man" from its natural state; i.e., that, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a naturally occurring polynucleotide naturally present in a living organism in its natural state is not "isolated," but the same polynucleotide separated from coexisting materials of its natural state is "isolated", as the term is employed herein. As part of or following isolation, such polynucleotides can be joined to other polynucleotides, such as DNA, for mutagenesis, to form fusion proteins, and for propagation or expression in a host, for instance. The isolated polynucleotides, alone or joined to other polynucleotides such as vectors, can be introduced into host cells, in culture or in whole organisms. Introduced into host cells in

culture or in whole organisms, such DNA still would be isolated, as the term is used herein, because they would not be in their naturally occurring form or environment. Similarly, the polynucleotides may occur in a composition, such as a media formulations, solutions for introduction of polynucleotides, for example, into cells, compositions or solutions for chemical or enzymatic reactions, for instance, which are not naturally occurring compositions, and, therein remain isolated polynucleotides within the meaning of that term as it is employed herein.

"Plasmids". Starting plasmids disclosed herein are either commercially available, publicly available, or can be constructed from available plasmids by routine application of well known, published procedures. Many plasmids and other cloning and expression vectors that can be used in accordance with the present invention are well known and readily available to those of skill in the art. Moreover, those of skill readily may construct any number of other plasmids suitable for use in the invention.

"Polynucleotides(s)" of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The DNA may be double-stranded or single-stranded. Single-stranded polynucleotides may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand. Polynucleotides generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among others, single-and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-,

double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded, or a mixture of single- and double-stranded regions. In addition, polynucleotide as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term polynucleotide also includes DNA or DNA that contain one or more modified bases. Thus, DNA or DNA with backbones modified for stability or for other reasons are "polynucleotides" as that term is intended herein. Moreover, DNA or DNA comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including simple and complex cells, inter alia. Polynucleotides embraces short polynucleotides often referred to as oligonucleotide(s). It will also be appreciated that RNA made by transcription of this doubled stranded nucleotide sequence, and an antisense strand of a nucleic acid molecule of the invention or an oligonucleotide fragment of the nucleic acid molecule, are contemplated within the scope of the invention. An antisense sequence is constructed by inverting the sequence of a nucleic acid molecule of the invention. relative to its normal presentation for transcription. Preferably, an antisense sequence is

constructed by inverting a region preceding the initiation codon or an unconserved region.

The antisense sequences may be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art.

"Stringent hybridization conditions" are those which are stringent enough to provide specificity, reduce the number of mismatches and yet are sufficiently flexible to allow formation of stable hybrids at an acceptable rate. Such conditions are known to those skilled in the art and are described, for example, in Sambrook, et al, (1989). By way of example only, stringent hybridization with short nucleotides may be carried out at 5-10° below the T_M using high concentrations of probe such as 0.01-1.0 pmole/ml. Preferably, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences.

"Variant(s)" of polynucleotides are polynucleotides that differ in nucleotide sequence from another, reference polynucleotide. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. Changes in the nucleotide sequence of the variant may be silent. That is, they may not alter the amino acids encoded by the polynucleotide. Where alterations are limited to silent changes of this type a variant will encode a polypeptide or polynucleotide with the same amino acid sequence as the reference. Changes in the nucleotide sequence of the variant may alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Such nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide or polynucleotide encoded by the reference sequence.

As hereinbefore mentioned, the present inventors have identified and sequenced a DNA sequence encoding PDE10A. The DNA sequence is shown in the Sequence Listing as SEQ ID NO:1, NO:2 and NO:11.

It will be appreciated that the invention includes nucleotide or amino acid sequences which have substantial sequence homology with the nucleotide sequences shown in the Sequence Listing as SEQ ID NO:1, NO:2 and NO:11. The term "sequences having substantial sequence homology" means those nucleotide and amino acid sequences which have slight or inconsequential sequence variations from the sequences disclosed in the Sequence Listing as SEQ ID NO:1, NO:2 and NO:11; i.e. the homologous sequences function in substantially the same manner to produce substantially the same polypeptides as the actual sequences. The variations may be attributable to local mutations or structural modifications. It is expected that a sequence having 85-90% sequence homology with the DNA sequence of the invention will provide a functional PDE10 polypeptide.

As used herein, "PDE10A" comprises a polynucleotide sequence which is down regulated in the course of CAG repeat disorders selected from the group consisting of: (a) a sequence comprising SEQ ID NO:1; (b) a sequence comprising SEQ ID NO:2; (c) a sequence comprising SEQ ID NO:11; (d) a sequence comprising nucleotides 257 to 2596 of SEQ ID NO:11; (e) a sequence which is at least 90% homologous with a sequence of (a), (b), (c) or (d), and; (f) a sequence which hybridizes to (a), (b), (c) or (d) under stringent conditions. In an embodiment, the isolated polynucleotide segment is cDNA. The invention also teaches an isolated polynucleotide segment, which retains substantially the same biological function or

activity as the polynucleotide encoded by the polynucleotide sequence.

Further embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding PDE10 polypeptide or polynucleotide, and polynucleotides which are complementary to such polynucleotides. Other embodiments are polynucleotides that comprise a region that is at least 80% identical over their entire length to a polynucleotide encoding PDE10 of SEQ ID NO.11 and polynucleotides complementary thereto. This includes polynucleotides at least 90% identical over their entire length to the same, and among these embodiments are polynucleotides with at least 95%. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

The polynucleotides of the present invention may be employed as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein.

Analysis of the complete nucleotide and amino acid sequences of the protein of the invention using the procedures of Sambrook et al., supra, have been used to determine the expressed region, initiation codon and untranslated sequences of the PDE10A gene. The transcription regulatory sequences of the gene are determined by analyzing fragments of the DNA for their ability to express a reporter gene such as the bacterial gene lacZ.

The nucleic acid molecules of the invention allow those skilled in the art to construct

nucleotide probes for use in the detection of nucleotide sequences in biological materials. As shown in FIG. 11, 13, 15 and 16, a number of unique restriction sequences for restriction enzymes are incorporated in the nucleic acid molecule identified in the Sequence Listing as SEQ ID NO:1, NO:2 and NO:11, and these provide access to nucleotide sequences which code for polypeptides unique to the PDE10A polypeptide of the invention. Nucleotide sequences unique to PDE10A or isoforms thereof, can also be constructed by chemical synthesis and enzymatic ligation reactions carried out by procedures known in the art.

A nucleotide probe may be labeled with a detectable marker such as a radioactive label which provides for an adequate signal and has sufficient half-life such as 32p, 3H, 14C or the like. Other detectable markers which may be used include antigens that are recognized by a specific labeled antibody, fluorescent compounds, enzymes, antibodies specific for a labeled antigen, and chemiluminescent compounds. An appropriate label may be selected having regard to the rate of hybridization and binding of the probe to the nucleotide to be detected and the amount of nucleotide available for hybridization. The nucleotide probes may be used to detect genes related to or analogous to PDE10A of the invention.

Accordingly, the present invention also provides a method of detecting the presence of nucleic acid molecules encoding a polypeptide related to or analogous to PDE10A in a sample comprising contacting the sample under hybridization conditions with one or more of the nucleotide probes of the invention labeled with a detectable marker, and determining the degree of hybridization between the nucleic acid molecule in the sample and the nucleotide probes.

Hybridization conditions which may be used in the method of the invention are known in the art and are described for example in Sambrook J, et al., *supra*. The hybridization product may be assayed using techniques known in the art. The nucleotide probe may be labeled with a detectable marker as described herein and the hybridization product may be assayed by detecting the detectable marker or the detectable change produced by the detectable marker.

The nucleic acid molecule of the invention also permits the identification and isolation, or synthesis of nucleotide sequences which may be used as primers to amplify a polynucleotide molecule of the invention, for example in polymerase chain reaction (PCR). The length and bases of the primers for use in the PCR are selected so that they will hybridize to different strands of the desired sequence and at relative positions along the sequence such that an extension product synthesized from one primer when it is separated from its template can serve as a template for extension of the other primer into a nucleic acid of defined length.

Primers which may be used in the invention are oligonucleotides i.e. molecules containing two or more deoxyribonucleotides of the nucleic acid molecule of the invention which occur naturally as in a purified restriction endonuclease digest or are produced synthetically using techniques known in the art such as, for example, phosphotriester and phosphodiester methods (See Good et al, 1977) or automated techniques (see, for example, Conolly, B. A., 1987). The primers are capable of acting as a point of initiation of synthesis when placed under conditions which permit the synthesis of a primer extension product which is complementary to the DNA sequence of the invention e.g. in the presence of nucleotide substrates, an agent for polymerization such as DNA polymerase and at suitable temperature and pH. Preferably, the primers are sequences that do not form secondary structures by base

pairing with other copies of the primer or sequences that form a hair pin configuration. The primer may be single or double-stranded. When the primer is double-stranded it may be treated to separate its strands before using it to prepare amplification products. The primer preferably contains between about 7 and 25 nucleotides.

The primers may be labeled with detectable markers which allow for detection of the amplified products. Suitable detectable markers are radioactive markers such as P-32, S-35, I-125, and H-3, luminescent markers such as chemiluminescent markers, preferably luminol, and fluorescent markers, preferably dansyl chloride, fluorcein-5-isothiocyanate, and 4-fluor-7-nitrobenz-2-axa-1,3 diazole, enzyme markers such as horseradish peroxidase, alkaline phosphatase, .beta.-galactosidase, acetylcholinesterase, or biotin.

It will be appreciated that the primers may contain non-complementary sequences provided that a sufficient amount of the primer contains a sequence which is complementary to a nucleic acid molecule of the invention or oligonucleotide sequence thereof, which is to be amplified. Restriction site linkers may also be incorporated into the primers allowing for digestion of the amplified products with the appropriate restriction enzymes facilitating cloning and sequencing of the amplified product.

Thus, a method of determining the presence of a nucleic acid molecule having a sequence encoding PDE10A or a predetermined oligonucleotide fragment thereof in a sample, is provided comprising treating the sample with primers which are capable of amplifying the nucleic acid molecule or the predetermined oligonucleotide fragment thereof in a polymerase chain reaction to form amplified sequences, under conditions which permit the formation of

amplified sequences and, assaying for amplified sequences.

The polymerase chain reaction refers to a process for amplifying a target nucleic acid sequence as generally described in Innis et al, Academic Press, 1989, in Mullis et al., U.S. Pat. No. 4,863,195 and Mullis, U.S. Pat. No. 4,683,202 which are incorporated herein by reference. Conditions for amplifying a nucleic acid template are described in M. A. Innis and D. H. Gelfand, 1989, which is also incorporated herein by reference.

The amplified products can be isolated and distinguished based on their respective sizes using techniques known in the art. For example, after amplification, the DNA sample can be separated on an agarose gel and visualized, after staining with ethidium bromide, under ultra violet (UV) light. DNA may be amplified to a desired level and a further extension reaction may be performed to incorporate nucleotide derivatives having detectable markers such as radioactive labeled or biotin labeled nucleoside triphosphates. The primers may also be labeled with detectable markers. The detectable markers may be analyzed by restriction and electrophoretic separation or other techniques known in the art.

The conditions which may be employed in the methods of the invention using PCR are those which permit hybridization and amplification reactions to proceed in the presence of DNA in a sample and appropriate complementary hybridization primers. Conditions suitable for the polymerase chain reaction are generally known in the art. For example, see M. A. Innis and D. H. Gelfand, 1989, which is incorporated herein by reference. Preferably, the PCR utilizes polymerase obtained from the thermophilic bacterium Thermus aquatics (Taq polymerase, GeneAmp Kit, Perkin Elmer Cetus) or other thermostable polymerase may be used to amplify

DNA template strands.

It will be appreciated that other techniques such as the Ligase Chain Reaction (LCR) and Nucleic-Acid Sequence Based Amplification (NASBA) may be used to amplify a nucleic acid molecule of the invention. In LCR, two primers which hybridize adjacent to each other on the target strand are ligated in the presence of the target strand to produce a complementary strand (Barney, 1991 and European Published Application No. 0320308, published Jun. 14, 1989). NASBA is a continuous amplification method using two primers, one incorporating a promoter sequence recognized by an RNA polymerase and the second derived from the complementary sequence of the target sequence to the first primer (U.S. Ser. No. 5,130,238 to Malek).

The present invention also teaches vectors which comprise a polynucleotide or polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of polynucleotides of the invention by recombinant techniques.

In accordance with this aspect of the invention the vector may be, for example, a plasmid vector, a single or double-stranded phage vector, a single or double-stranded RNA or DNA viral vector. In certain embodiments in this regard, the vectors provide for specific expression. Such specific expression may be inducible expression or expression only in certain types of cells or both inducible and cell-specific. Particular among inducible vectors are vectors that can be induced for expression by environmental factors that are easy to manipulate, such as temperature and nutrient additives. A variety of vectors suitable to this

aspect of the invention, including constitutive and inducible expression vectors for use in prokaryotic and eukaryotic hosts, are well known and employed routinely by those of skill in the art. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids, all may be used for expression in accordance with this aspect of the present invention.

The following vectors, which are commercially available, are provided by way of example. Among vectors for use in bacteria are pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia, and pBR322 (ATCC 37017). Among eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. These vectors are listed solely by way of illustration of the many commercially available and well known vectors that are available to those of skill in the art for use in accordance with this aspect of the present invention. It will be appreciated that any other plasmid or vector suitable for, for example, introduction, maintenance, propagation or expression of a polynucleotide or polypeptide of the invention in a host may be used in this aspect of the invention. Generally, any vector suitable to maintain, propagate or express polynucleotides to express a polypeptide or polynucleotide in a host may be used

for expression in this regard.

The appropriate DNA sequence may be inserted into the vector by any of a variety of well-known and routine techniques. In general, expression constructs will contain sites for transcription initiation and termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will include a translation initiating AUG at the beginning and a termination codon appropriately positioned at the end of the polynucleotide to be translated.

The DNA sequence in the expression vector is operatively linked to appropriate expression control sequence(s), including, for instance, a promoter to direct mRNA transcription.

Promoter regions can be selected from any desired gene using vectors that contain a reporter transcription unit lacking a promoter region, such as a chloramphenicol acetyl transferase ("CAT") transcription unit, downstream of restriction site or sites for introducing a candidate promoter fragment; i.e., a fragment that may contain a promoter. As is well known, introduction into the vector of a promoter-containing fragment at the restriction site upstream of the cat gene engenders production of CAT activity, which can be detected by standard CAT assays. Vectors suitable to this end are well known and readily available, such as pKK232-8 and pCM7. Promoters for expression of polynucleotides of the present invention include not only well known and readily available promoters, but also promoters that readily may be obtained by the foregoing technique, using a reporter gene. Among known prokaryotic promoters suitable for expression of polynucleotides and polypeptides in accordance with the present invention are the E. coli lacI and lacZ and promoters, the T3 and T7 promoters, the gpt promoter, the lambda PR, PL promoters and the trp promoter. Among

known eukaryotic promoters suitable in this regard are the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus ("RSV"), and metallothionein promoters, such as the mouse metallothionein-I promoter.

Vectors for propagation and expression generally will include selectable markers and amplification regions, such as, for example, those set forth in Sambrook et al., supra.

As hereinbefore mentioned, the present invention also teaches host cells which are genetically engineered with vectors of the invention.

Polynucleotide constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. The PDE10A polynucleotide or polypeptide products or isoforms or parts thereof, may be obtained by expression in a suitable host cell using techniques known in the art. Suitable host cells include prokaryotic or eukaryotic organisms or cell lines, for example bacterial, mammalian, yeast, or other fungi, viral, plant or insect cells. Methods for transforming or transfecting cells to express foreign DNA are well known in the art (See for example, Itakura et al., U.S. Pat. No. 4,704,362; Hinnen et al., 1978; Murray et al., U.S. Pat. No. 4,801,542; Upshall et al., U.S. Pat. No. 4,935,349; Hagen et al., U.S. Pat. No. 4,784,950; Axel et al., U.S. Pat. No. 4,399,216; Goeddal et al., U.S. Pat. No. 4,766,075; and Sambrook et al, 1989, all of which are incorporated herein by reference). Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2

and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

Host cells can be genetically engineered to incorporate polynucleotides and express polynucleotides of the present invention. Introduction of polynucleotides into the host cell can be affected by calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al. (1986) and Sambrook et al. (1989).

As hereinbefore mentioned, the present invention also teaches the production of polynucleotides of the invention by recombinant techniques.

The PDE10 polynucleotides encode a polypeptide which is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case in vivo, the additional amino acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed

such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In sum, a polynucleotide of the present invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences which are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

The polypeptides of the invention may be prepared by culturing the host/vector systems described above, in order to express the recombinant polypeptides. Recombinantly produced PDE10A based protein or parts thereof, may be further purified using techniques known in the art such as commercially available protein concentration systems, by salting out the protein followed by dialysis, by affinity chromatography, or using anion or cation exchange resins.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using DNA derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook et al., supra.

Polynucleotides of the invention, encoding the heterologous structural sequence of a

polynucleotide or polypeptide of the invention generally will be inserted into a vector using standard techniques so that it is operably linked to the promoter for expression. The polynucleotide will be positioned so that the transcription start site is located appropriately 5' to a ribosome binding site. The ribosome binding site will be 5' to the AUG that initiates translation of the polynucleotide or polypeptide to be expressed. Generally, there will be no other open reading frames that begin with an initiation codon, usually AUG, and lie between the ribosome binding site and the initiation codon. Also, generally, there will be a translation stop codon at the end of the expressed polynucleotide and there will be a polyadenylation signal in constructs for use in eukaryotic hosts. Transcription termination signal appropriately disposed at the 3' end of the transcribed region may also be included in the polynucleotide construct.

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polynucleotide or polypeptide. These signals may be endogenous to the polynucleotide or they may be heterologous signals. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well know to those skilled in the art. PDE10A polynucleotide or polypeptide can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for

purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polynucleotide is denatured during isolation and or purification.

In an embodiment, a nucleic acid molecule of the invention may be cloned into a glutathione S-transferase (GST) gene fusion system for example the pGEX-1 T, pGEX-2T and pGEX-3X of Pharmacia. The fused gene may contain a strong lac promoter, inducible to a high level of expression by IPTG, as a regulatory element. Thrombin or factor Xa cleavage sites may be present which allow proteolytic cleavage of the desired polypeptide from the fusion product. The glutathione S-transferase-PDE10A fusion protein may be easily purified using a glutathione sepharose 4B column, for example from Pharmacia. The 26 kd glutathione S-transferase polypeptide can be cleaved by thrombin (pGEX-1 or pGEX-2T) or factor Xa (pGEX-3X) and resolved from the using the polypeptide using the same affinity column. Additional chromatographic steps can be included if necessary, for example Sephadex or DEAE cellulose. The two enzymes may be monitored by protein and enzymatic assays and purity may be confirmed using SDS-PAGE.

The PDE10A protein or parts thereof may also be prepared by chemical synthesis using techniques well known in the chemistry of proteins such as solid phase synthesis (Merrifield, 1964) or synthesis in homogenous solution (Houbenweyl, 1987).

Within the context of the present invention, PDE10A polypeptide includes various structural forms of the primary protein which retain biological activity. For example, PDE10A polypeptide may be in the form of acidic or basic salts or in neutral form. In addition,

individual amino acid residues may be modified by oxidation or reduction. Furthermore, various substitutions, deletions or additions may be made to the amino acid or nucleic acid sequences, the net effect being that biological activity of PDE10A is retained. Due to code degeneracy, for example, there may be considerable variation in nucleotide sequences encoding the same amino acid.

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals but also additional heterologous functional regions. Thus, for instance, a region of additional amino acids, particularly charged amino acids, may be added to the C- or N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification or during subsequent handling and storage. Also, fusion proteins may be added to the polynucleotide or polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polynucleotide or polypeptide. The addition of peptide moieties to polynucleotide or polypeptides to engender secretion or excretion, to improve stability or to facilitate purification, among others, are familiar and routine techniques in the art. In drug discovery, for example, proteins have been fused with antibody Fc portions for the purpose of high-throughput screening assays to identify antagonists (see Bennett et al., 1995, and Johanson et al., 1995).

Detecting Presence of or Predisposition for CAG Repeat Disorders

This invention is also related to the use of the PDE10A polynucleotides to detect complementary polynucleotides as a diagnostic reagent. Detection of the level of expression of PDE10A in a eukaryote, particularly a mammal, and especially a human, will provide a

method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, exhibiting decreased levels of PDE10A may be detected by a variety of techniques. Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as the striatum, nucleus accumbens and olfactory tubercule. RNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki et al., 1986) prior to analysis. As an example, PCR primers complementary to the nucleic acid encoding PDE10A can be used to identify and analyze PDE10A presence and/or expression. Using PCR, characterization of the level of PDE10A present in the individual may be made by comparative analysis.

The invention thus provides a process for detecting disease by using methods known in the art and methods described herein to detect decreased expression of PDE10 polynucleotide. For example, decreased expression of PDE10 polynucleotide can be measured using any on of the methods well known in the art for the quantification of polynucleotides, such as, for example, PCR, RT-PCR, DNAse protection, northern blotting and other hybridization methods. Thus, the present invention provides a method for detecting triplet-repeat disorders, and a method for detecting a genetic pre-disposition for triplet-repeat disorders and other disorders of the basal ganglia including schizophrenia, stroke, trauma, Parkinson's disease and Alzheimer's disease (AD). More generally, the present invention provides a method for detecting a genetic pre-disposition for neurological disorders characterized by progressive cell loss.

Drug Screening Assays

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of PDE10 polypeptides or polynucleotides, such as its interaction with PDE10-binding molecules. The identification of mutations in specific genes in inherited neurodegenerative disorders, combined with advances in the field of transgenic methods, provides those of skill in the art with the information necessary to further study human diseases. This is extraordinarily useful in modeling familial forms of triplet-repeat disorders and other disorders of the basal ganglia including schizophrenia, stroke, trauma, Parkinson's disease and Alzheimer's disease (AD). More generally, the present invention is useful for modeling neurological disorders characterized by progressive cell loss, as well as those involving acute cell loss, such as stroke and trauma.

For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, may be prepared from a cell that expresses a molecule that binds PDE10. The preparation is incubated with labeled PDE10 in the absence or the presence of a candidate molecule which may be a PDE10 agonist or antagonist. The ability of the candidate molecule to bind the binding molecule is reflected in decreased binding of the labeled ligand.

PDE10-like effects of potential agonists and antagonists may by measured, for instance, by determining activity of a reporter system following interaction of the candidate molecule with a cell or appropriate cell preparation, and comparing the effect with that of PDE10 or molecules that elicit the same effects as PDE10. Reporter systems that may be useful in this

regard include, but are not limited to, colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in PDE10 activity, and binding assays known in the art.

Another example of an assay for PDE10 antagonists is a competitive assay that combines PDE10 and a potential antagonist with membrane-bound PDE10-binding molecules, recombinant PDE10 binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. PDE10 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of PDE10 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing PDE10-induced activities, thereby preventing the action of PDE10 by excluding PDE10 from binding.

Potential antagonists include a small molecule which binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, 1988, for a description of these molecules).

Potential antagonists include compounds related to and derivatives of PDE10.

Developing modulators of the biological activities of specific PDEs requires differentiating PDE isozymes present in a particular assay preparation. The classical enzymological approach of isolating PDEs from natural tissue sources and studying each new isozyme may be used. Another approach has been to identify assay conditions which might favor the contribution of one isozyme and minimize the contribution of others in a preparation. Still another approach has been the separation of PDEs by immunological means. Each of the foregoing approaches for differentiating PDE isozymes is time consuming. As a result many attempts to develop selective PDE modulators have been performed with preparations containing more than one isozyme. Moreover, PDE preparations from natural tissue sources are susceptible to limited proteolysis and may contain mixtures of active proteolytic products that have different kinetic, regulatory and physiological properties than the full length PDEs.

Recombinant PDE10 polypeptide products of the invention greatly facilitate the development of new and specific PDE10 modulators. The need for purification of an isozyme can be avoided by expressing it recombinantly in a host cell that lacks endogenous phosphodiesterase activity (e.g., yeast strain YKS45 deposited as ATCC 74225). Once a compound that modulates the activity of the PDE10 is discovered, its selectivity can be evaluated by comparing its activity on the PDE10 to its activity on other PDE isozymes. Thus, the combination of the recombinant PDE10 products of the invention with other recombinant PDE products in a series of independent assays provides a system for developing selective modulators of PDE10. Selective modulators may include, for example, antibodies and other proteins or peptides which specifically bind to the PDE10 or PDE10 nucleic acid,

oligonucleotides which specifically bind to the PDE10 (see Patent Cooperation Treaty International Publication No. WO93/05182 published Mar. 18, 1993 which describes methods for selecting oligonucleotides which selectively bind to target biomolecules) or PDE10 nucleic acid (e.g., antisense oligonucleotides) and other non-peptide natural or synthetic compounds which specifically bind to the PDE10 or PDE10 nucleic acid. Mutant forms of the PDE10 which alter the enzymatic activity of the PDE10 or its localization in a cell are also contemplated. Crystallization of recombinant PDE10 alone and bound to a modulator, analysis of atomic structure by X-ray crystallography, and computer modelling of those structures are methods useful for designing and optimizing non-peptide selective modulators. See, for example, Erickson et al., *Ann. Rep. Med. Chem.*, 27: 271-289 (1992) for a general review of structure-based drug design.

Targets for the development of selective modulators include, for example: (1) the regions of the PDE10 which contact other proteins and/or localize the PDE10 within a cell, (2) the regions of the PDE10 which bind substrate, (3) the allosteric cGMP-binding site(s) of PDE10, (4) the metal-binding regions of the PDE10, (5) the phosphorylation site(s) of PDE10 and (6) the regions of the PDE10 which are involved in dimerization of PDE10 subunits.

Thus, the present invention provides a method for screening and selecting compounds which promote triplet-repeat disorders, and a method for screening and selecting compounds which treat or inhibit triplet-repeat disorders, as well as schizophrenia, stroke, trauma, Parkinson's disease and Alzheimer's disease. More generally, the present invention provides a method for screening and selecting compounds which promote or inhibit neurological disorders characterized by progressive cell loss, as well as those involving acute cell loss, such as

stroke and trauma.

The selected antagonists and agonists may be administered, for instance, to inhibit progressive and acute neurological disorders, such as Huntington's disease, Parkinson's disease, schizophrenia, Alzheimer's disease (AD), stroke or trauma.

Antagonists and agonists and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by direct microinjection into the affected area, or by intravenous or other routes. These compositions of the present invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of antagonists or agonists of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation is prepared to suit the mode of administration.

Inhibition of PDE10A will be highly detrimental to striatal brain function. The progressive decline in PDE10A mRNA levels in HD may lead to dysregulation of cAMP levels and neuronal dysfunction. Up-regulation of PDE10A will be effective in combating such neuronal dysfunction.

Gene Therapy

A variety of gene therapy approaches may be used in accordance with the invention to modulate expression of the PDE10A gene in vivo. For example, antisense DNA molecules may be engineered and used to block translation of PDE10A mRNA in vivo. Alternatively, ribozyme molecules may be designed to cleave and destroy the PDE10A mRNAs in vivo. In another alternative, oligonucleotides designed to hybridize to the 5' region of the PDE10A gene (including the region upstream of the coding sequence) and form triple helix structures may be used to block or reduce transcription of the PDE10A gene. In yet another alternative, nucleic acid encoding the full length wild-type PDE10A message may be introduced in vivo into cells which otherwise would be unable to produce the wild-type PDE10A gene product in sufficient quantities or at all.

In a preferred embodiment, the antisense, ribozyme and triple helix nucleotides are designed to inhibit the translation or transcription of PDE10A. To accomplish this, the oligonucleotides used should be designed on the basis of relevant sequences unique to PDE10A.

For example, and not by way of limitation, the oligonucleotides should not fall within those region where the nucleotide sequence of PDE10A is most homologous to that of other PDEs, such as PDE2 PDE5 and PDE6, herein referred to as "unique regions".

In the case of antisense molecules, it is preferred that the sequence be chosen from the unique regions. It is also preferred that the sequence be at least 18 nucleotides in length in order to

achieve sufficiently strong annealing to the target mRNA sequence to prevent translation of the sequence. Izant and Weintraub, 1984, Cell, 36:1007-1015; Rosenberg et al., 1985, Nature, 313:703-706.

In the case of the "hammerhead" type of ribozymes, it is also preferred that the target sequences of the ribozymes be chosen from the unique regions. Ribozymes are RNA molecules which possess highly specific endoribonuclease activity. Hammerhead ribozymes comprise a hybridizing region which is complementary in nucleotide sequence to at least part of the target RNA, and a catalytic region which is adapted to cleave the target RNA. The hybridizing region contains nine (9) or more nucleotides. Therefore, the hammerhead ribozymes of the present invention have a hybridizing region which is complementary to the sequences listed above and is at least nine nucleotides in length. The construction and production of such ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, 1988, Nature, 334:585-591.

The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in Tetrahymena Thermophila (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, Science, 224:574-578; Zaug and Cech, 1986, Science, 231:470-475; Zaug, et al., 1986, Nature, 324:429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, Cell, 47:207-216). The Cech endoribonucleases have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences

that are present in PDE10A but not other PDEs.

The foregoing compounds can be administered by a variety of methods which are known in the art including, but not limited to the use of liposomes as a delivery vehicle. Naked DNA or RNA molecules may also be used where they are in a form which is resistant to degradation such as by modification of the ends, by the formation of circular molecules, or by the use of alternate bonds including phosphothionate and thiophosphoryl modified bonds. In addition, the delivery of nucleic acid may be by facilitated transport where the nucleic acid molecules are conjugated to poly-lysine or transferrin. Nucleic acid may also be transported into cells by any of the various viral carriers, including but not limited to, retrovirus, vaccinia, AAV, and adenovirus.

Alternatively, a recombinant nucleic acid molecule which encodes, or is, such antisense, ribozyme, triple helix, or PDE10A molecule can be constructed. This nucleic acid molecule may be either RNA or DNA. If the nucleic acid encodes an RNA, it is preferred that the sequence be operatively attached to a regulatory element so that sufficient copies of the desired RNA product are produced. The regulatory element may permit either constitutive or regulated transcription of the sequence. In vivo, that is, within the cells or cells of an organism, a transfer vector such as a bacterial plasmid or viral RNA or DNA, encoding one or more of the RNAs, may be transfected into cells e.g. (Llewellyn et al., 1987, J. *Mol. Biol.*, 195:115-123; Hanahan et al. 1983, *J. Mol. Biol.*, 166:557-580). Once inside the cell, the transfer vector may replicate, and be transcribed by cellular polymerases to produce the RNA or it may be integrated into the genome of the host cell. Alternatively, a transfer vector containing sequences encoding one or more of the RNAs may be transfected into cells or

introduced into cells by way of micromanipulation techniques such as microinjection, such that the transfer vector or a part thereof becomes integrated into the genome of the host cell.

Composition, Formulation, and Administration of Pharmaceutical Compositions

The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules,

liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In

addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds

may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer,

and an aqueous phase. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Many of the compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents that are the corresponding

free base forms.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, transdermal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into an affected area, often in a depot or sustained release formulation.

Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with an antibody specific for affected cells. The liposomes will be targeted to and taken up selectively by the cells.

The pharmaceutical compositions generally are administered in an amount effective for treatment or prophylaxis of a specific indication or indications. It is appreciated that optimum dosage will be determined by standard methods for each treatment modality and indication, taking into account the indication, its severity, route of administration, complicating conditions and the like. In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms associated with such disorders. Techniques for formulation and administration of the compounds of the instant

application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition. For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.001 mg/kg to 10 mg/kg, typically around 0.01 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

The invention further provides diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, reflecting approval by the agency of the manufacture, use or sale of the product for human administration.

EXAMPLES

The present invention is further described by the following examples. These examples, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

EXAMPLE 1 - Isolation of PDE10A

Wild-type (B6CBAF1) and HD transgenic [B6CBA-TgN(Hdexon1)62Gpb] mice (Jackson Laboratories) and adult Sprague-Dawley rats (250-300 g; Charles River Laboratories) and were used in this study. The genotype of the mice was determined by PCR amplification of a 100 bp region of the integrated human HD exon 1 transgene using primers corresponding to nts 3340-3459 (5'-AGG GCT GTC AAT CAT GCT GG-3') and nts 3836-3855 (5'-AAA CTC ACG GTC GGT GCA GC-3') of clone E4.1 of the human HD gene (Accession number L34020). PCR conditions used are described in Mangiarini et al.(1996). DNA was extracted from a tail clip and an ear punch from each mouse used in this study. Both samples were subjected to PCR genotype analysis. For *in situ* hybridization analysis, the animals were anesthetized with >100 mg/kg sodium pentobarbital, decapitated, the brains removed and stored at -70°C prior to sectioning. For RNA isolation, animals were anesthetized, decapitated and the striatum and cortex were excised and stored in liquid nitrogen prior to RNA extraction. Animal care was given according to protocols approved by Dalhousie University and the Canadian Council of Animal Care.

Differential display was used to identify novel mDNA or previously described mDNA whose relative expression levels are altered as a result of the presence of the transgene. Using differential display, the mRNA populations derived from the striatum of 10 week old wild type were compared with age-matched R6/2 transgenic mice. Differential display has been used extensively (> 750 references) since its development (Liang and Pardee, 1992) to identify changes in gene expression in cells and in tissues including brain (Douglass et al., 1995; Babity et al., 1997a; Livesey et al., 1997; Berke et al., 1998). Perhaps the most

important finding was the demonstration by Qu et al., (1996) that differential display can be used to isolate genes differentially expressed in inbred strains of mice. The power of differential display is that the sequence information obtained can be directly related to the experimental paradigm. Moreover, such sequence information includes sufficient information to identify transcripts and can then lead to experiments that reveal function of the cognate protein in the experimental model.

DNA sequence information of potentially differentially expressed cDNA can be used to generate oligonucleotide probes for in situ hybridization to define the anatomical and temporal patterns of expression of specific transcripts (see Babity et al., 1997a). This technique is especially useful to study changes in steady-state levels of mRNA in heterogeneous tissue such as brain. Brain tissue can be micro-dissected (Babity et al., 1997b). This enabled the present inventors to reduce the requirement for tissue, and hence compare the mRNA populations derived from individual animals for each experimental group.

Thus RT-PCR (Denovan-Wright et al., 1999) was used to identify differences in the patterns of gene expression between the striatum of wild-type and transgenic mice that were hemizygous for the 5' UTR, exon 1 and part of intron 1 of the human Huntingon's Disease gene. Total cellular RNA was isolated from the striatum and cortex of three 10 week-old wild-type and three 10 week-old R6/2 HD mice (Mangiarini et al., 1996) and used as the template to generate single-stranded cDNA. Total cellular RNA from each animal and tissue was purified using TrizolTM reagent (Gibco BRL) and the manufacture's protocol. 10 μg aliquots of total RNA were treated with RQ1 DNAse-free DNAse (Promega) in the presence

of DNAsin™ (Promega) DNAse inhibitor to remove trace genomic DNA and then converted to single-stranded cDNA. The primers and conditions for PCR amplification follow those of the Delta™ RNA fingerprinting manual (Clontech).

The cDNA was then used as the substrate for PCR reactions using 57 differential display primer combinations. The radio-labelled PCR products were fractionated on a denaturing acrylamide sequencing gels using a Genomyx LR™ sequencing apparatus, transferred to 3MM filter paper and dried. The dried acrylamide gels were exposed to autoradiography film (BioMax MR™) overnight. After fractionating the radio-labelled PCR products on denaturing acrylamide gels, it was found that the overwhelming majority of the approximately 18,000 PCR products screened were common to both the wild-type and HD mice (data not shown). One PCR product, amplified using the primers P7 (5'-ATT AAC CCT CAC TAA ATG CTG TAT G-3') and T6 (5'-CAT TAT GCT GAG TGA TAT CTT TTT TTT TCG-3') of approximately 500 bp, was observed in each of three samples derived from the striatum of wild-type mice (FIG. 1). This 500 bp band was absent from the samples derived from the striatum of the HD mice (FIG. 1) and was absent from each of the samples derived from the cortical tissue (data not shown).

FIG. 1 shows the Down-regulated in Huntington's Disease (PDE10A) transcript, identified by differential display RT PCR. A band of approximately 500 bp (arrow) was amplified from cDNA made form 10 week-old wild-type but not 10 week-old HD striatal tissue. Total RNA from individual animals (numbered 1-6) was used as the substrate for the generation of single-stranded cDNA. Animals 1, 2 and 3 were transgenic HD mice. Animals 4, 5 and 6 were wild-type mice.

EXAMPLE 2 - Cloning of PDE10A

The 500 bp band, designate PDE10Apcr, was excised from the dried gel and rehydrated in 40 μl of H₂O for 10 min at room temperature. The eluted DNA was subjected to PCR reamplification using the P7 and T6 primers, rTaq polymerase (Pharmacia) and the following conditions: 60" @ 94°C, 19 x (30" @ 94°C, 30" @ 58°C, 120" @ 68°C + 4" per cycle), 7' @ 68°C. The PCR reaction was subjected to agarose gel electrophoresis and the 500 bp band was removed from the gel, extracted from the agarose using the Qiagen gel extraction protocol and cloned into the vector, pGem-T using standard methods. Plasmid DNA was isolated from selected transformants using Qiagen spin columns. The resultant clone was named pPDE10A.

EXAMPLE 3 - Identification of PDE10A

The cloned insert of pPDE10A was radio-labelled and used as a hybridization probe in northern blot analysis (FIG. 2). Northern blots of total RNA were prepared using the method described in Denovan-Wright et al. (1998). The 500 bp cloned insert of PDE10A was radio-labelled with [α-32P]dCTP (3000 Ci/mmol) using the Ready-to-Go dCTP beads (Pharmacia). Northern blot hybridization, brain tissue preparation and *in situ* hybridization are described in Denovan-Wright et al. (1998). The 500 bp cloned insert of pPDE10A annealed to a transcript of approximately 9.5 kb in total RNA isolated from the striatum of ten week-old wild-type mice.

FIG. 2 demonstrates that PDE10A is expressed in the striatum but not the cortex of wild-type mice and the steady-state levels of PDE10A are reduced in 10 week old transgenic HD mice. The differential expression of PDE10A in HD mice was confirmed by northern blot analysis. The cloned insert of pPDE10A was radio-labelled and used as a hybridization probe in northern blot analysis. The northern blot was prepared by size-fractionating total RNA from the striatum and cortex of three individual 10 week-old HD (1, 2 and 3) and wild-type (4, 5 and 6) mice. Following the hybridization of pPDE10A, the radio-label was removed and the blot was subsequently allowed to hybridize with a probe that detects constitutively expressed cyclophilin. The hybridization pattern of the cyclophilin probe is aligned below the northern blot demonstrating that equivalent amount of RNA were present in each lane. The relative mobility of RNA molecular weight standards (RNA ladder, Gibco BRL) are shown on the left of the northern blot.

The hybridization signal of pPDE10A was significantly lower in the RNA samples derived from the striatum of 10 week-old HD mice. No expression of the PDE10A mRNA was detected in the cortical RNA samples derived from either the wild-type or HD mice.

EXAMPLE 4 - Sequencing PDE10A

The sequence of the cloned differential display band, pPDE10A, was determined using M13 universal forward and reverse sequencing primers and the T7 sequencing kit (Pharmacia).

The 484 bp cDNA fragment did not have sequence similarity to any Genbank entries.

FIG. 3 shows the nucleotide sequence of the cloned PDE10A differential display product,

pPDE10A. The position of the primers used to amplify the fragment are underlined and labelled. The nucleotide sequence and position of oligonucleotide probes 1 and 2 within the pPDE10A sequence are shown.

EXAMPLE 5 - Isolation and Characterization of cDNA PDE10A

In order to isolate PDE10A cDNA clones, oligonucleotide probes 1 and 2 were used in 5' and 3' Rapid Amplification of cDNA Ends (RACE) reactions using commercially prepared RACE-ready mouse striatal cDNA (Clontech). Several independent clones were isolated and those that contained the sequence of pPDE10A were selected for further analysis. Each of the 5' RACE clones was identical in sequence over the length that the clones could be aligned. The difference in length between these clones is a result of termination of the original reverse-transcriptase reaction at different positions along the mRNA. No difference in size or sequence was detected between several 3' RACE clones. The longest 5' RACE clone and one 3' RACE clone were completely sequenced using internal primers. The present inventors were able to isolate a very short clone that extended the 5' RACE clone using an internal primer (probe 3, 5'- CTA TTT CAC AAG AGA CTG ACC AGC CAA TAA ATC TC- 3'). The compiled sequence of the first PDE10A cDNA clone, named cPDE10A-1 is presented in FIG. 10. cPDE10A-1 is 3235 bp in length. The restriction map of cPDE10A-1 is shown in FIG. 11.

The mRNA that hybridized with pPDE10A was approximately 9.5 kilobases in length. In order to obtain PDE10A cDNA clone that was larger than cPDE10-1, the present inventors screened a mouse brain cDNA library. Several clones were identified that hybridized with

the pPDE10 probe. The sequence of the largest of these cDNA clones, cPDE10-2, was determined. The sequence (FIG. 12) was 5753 base pairs in length. The restriction map of cPDE10-2 is shown in FIG. 13.

cPDE10-1 and cPDE10-2 share sequence identity over 2095 bp. However, the 5' 1142 bp of cPDE10-1 and the 5' 1689 bp of cPDE10-2 are unique to each clone. Clone cPDE10-2 extends 1969 bp in the 3' direction compared to cPDE10-1. A schematic showing the regions of sequence identity and the unique sequences of cPDE10-1 and -2 are shown in FIG. 14.

The compiled sequence of the mouse PDE10 cDNA clone, named cPDE10A, is presented in FIG. 15 with RACEs. A further sequence, without RACEs, is shown in FIG. 19. The coding sequence and restriction map of cPDE10A is shown in FIG. 16, and updated at FIG. 17. FIG. 18 is a restriction map of PDE10A. The coding region has a met initiator commencing at nucleotide 257, with a stop codon ending at nucleotide 2596.

PDE10A was found to have extremely high homology with human PDE10s identified by Loughney et al., WO99/42596, the contents of which are incorporated herein by reference.

EXAMPLE 6 - Localization of PDE10A in the Brain

In order to identify the coding strand and to localize the transcript in the wild-type mouse brain, two oligonucleotide probes were designed (probe 1, 5'- GAA CAT GTA GCA TAT ACT CCA GAC AAC AGA TCA TAT GG – 3'; probe 2, 5' – CAG CTT CTC CAC AGG AAC ACA GTA ACA AAG AG –3') that were complementary to different regions and

strands of the 484 bp pPDE10A clone. These oligonucleotides were used for *in situ* hybridization analysis. Using high stringency post *in situ* hybridization washes (2 x 30' in 1X SSC @ 58°C, 4 x 15' in 1X SSC @ 58°C, 4 x 15' in 0.5X SSC @ 58°C, 4 x 15' in 0.25X SSC @ 58°C), it was found that oligonucleotide probe 1 annealed with mRNA in the striatum, nucleus accumbens and olfactory tubercule of ten week-old wild-type mice (FIG. 4). The hybridization signal was significantly reduced in the striatum, nucleus accumbens and olfactory tubercle of the 10 week-old HD mice (FIG. 4).

FIG. 5 shows *in situ* hybridization of probe 1 to coronal (top three sections) and saggital (bottom section) 10 week-old wild-type (WT) and HD mouse brain sections. Specific hybridization of the probe was observed in the striatum, nucleus accumbens and olfactory tubercle of wild-type mice. The top three sections represent the distribution of PDE10A throughout the rostral-caudal axis of the striatum.

The *in situ* hybridization results confirmed the northern blot analysis demonstrating, 1) that the expression of PDE10A mRNA was restricted to the striatum, nucleus accumbens and olfactory tubercle and 2) that the levels of PDE10A mRNA were decreased in HD mice compared to the wild-type. The probe did not anneal with mRNA in any other brain nuclei. No hybridization of oligonucleotide probe 2 was observed in any region of the brain in wild-type or HD mice (Fig. 3). Based on this hybridization, the coding strand, complementary to probe 1, of pPDE10A was defined.

EXAMPLE 7 - Characterization of PDE10

The *in situ* hybridization using oligonucleotide probe 1 demonstrated that PDE10A mRNA levels in the striatum, nucleus accumbens and olfactory tubercule were decreased in ten week- old HD mice. By ten weeks of age, the HD mice all showed motor symptoms including resting tremor and stereotypic involuntary movements. Moreover, these mice immediately clasped their feet together and curled into a tight ball when picked up by their tails.

As the phenotypic signs are progressive over a number of weeks, the present inventors examined whether the PDE10A transcript was ever expressed in the striatum of the HD mice or whether the steady-state levels of the transcript diminished in the striatum in a course that parallelled the development of the motor disorders. Wild-type and HD mice were sacrificed at 5, 7 and 8 weeks of age and their brains were prepared for *in situ* hybridization analysis using probe 1 (FIG. 5).

FIG. 5 shows the levels of PDE10A mRNA decrease in HD mice over the period of time that the HD mice develop abnormal movements and postures. *In situ* hybridization analysis of coronal and saggital sections of wild-type and HD mouse brain using oligonucleotide probe 1 which is complementary to the coding strand of PDE10A. At 5 weeks of age, before the development of motor symptoms, the HD mice express the PDE10A transcript in the same brain nuclei and at the same relative levels as wild-type mice. The steady-state level PDE10A decreases in the striatum, nucleus accumbens and olfactory tubercle from 5 to 10 weeks in the HD but not wild-type mice. By 9 weeks of age, the HD mice have abnormal

movement and posture. The numbers refer to the age in weeks of the wild-type (WT) and Huntington's (HD) transgenic mice.

None of the mice at these ages had overt motor symptoms. Sections taken throughout the rostral-caudal axis of the striatum showed that PDE10A was expressed in the 5 week-old wild-type and HD mice. The relative hybridization of probe 1 did not change in 5, 7, 8 and 10 week-old wild-type mice. The intensity of the hybridization signal appeared to decrease in the striatum, nucleus accumbens and olfactory tubercle of HD mice from 5 to 10 weeks compared to their wild-type litter mates (FIG. 5).

The levels of PDE10A were significantly reduced by 8 weeks of age in the HD mice, using two in situ oligonucleotide probes, one complementary to the 3' UTR, the second complementary to an internal portion of the coding region. The hybridization pattern observed in the wild-type and HD mice was the same for both the probes employed. This analysis demonstrated that there is a reduction in the complete PDE10A mRNA levels during the development of the HD phenotype and not that there was a differential reduction in the PDE10A coding region as compared to the extensive 3' UTR. Moreover, *in situ* hybridization using the PDE10A-specific probe against neurologically normal and HD human brain tissue demonstrated that there was a decrease in PDE10A levels in human HD patients.

One day old wild-type and HD mice were frozen, sectioned on a cryostat and whole mouse sections were prepared for *in situ* hybridization using probe 1. The same high stringency post-hybridization washing conditions were employed for the one day-old mouse body sections as were used for the adult mouse brain sections. Parallel *in situ* hyridization

experiments using the probe 2 were performed in order to determine the level of non-specific signal in the mouse sections. Probe 1 specifically annealed to the developing striatum (FIG. 6).

FIG. 6 demonstrates that PDE10A is expressed in the developing striatum of one day-old wild-type and HD mice. The sections on the left were subjected to *in situ* hybridization using probe 1. Following hybridization, the sections were counter-stained with cresyl violet to visualize the mouse organs. The signal outside the brain was non-specific as probe 2 and other unrelated control oligonucleotide probes all labelled these tissues.

There was no difference in the pattern of hybridization between the one day-old wild-type and HD mice demonstrating that PDE10A was expressed in the developing brain of both wild-type and HD mice.

Following *in situ* hybridization, the sections were covered in autoradiographic emulsion, left in the dark to expose for 4 weeks and then developed and viewed under dark-field microscopy or, after counter-staining the sections with cresyl violet to visualize neuronal cell bodies, under bright-field microscopy. Silver grains were observed to be concentrated in the striatum of the wild-type mice. FIG. 7 shows emulsion autoradiography of mouse brain sections following *in situ* hybridization of probe 1 demonstrated that the PDE10A transcript is expressed in neurons. PDE10A is not homogeneously distributed throughout the mouse striatum. Dark field illumination of the sections after emulsion autoradiography showed that the silver grains were clustered in specific regions of the 10 week old wild-type mouse striatum (A and C). Sections from 10 week old HD mice subjected to identical *in situ* and

emulsion autoradiographic conditions are shown in B and D. The photomicrographs shown in A and B were viewed using the 10X objective (bar represents $100~\mu m$). The micrographs shown in C and D, were viewed under the 20X objective (bar represents $25~\mu m$). The insert in panel C is a portion of the section in A and C counter-stained with cresyl violet to visualize the neurons, viewed using the 40X objective under bright filed illumination. Note the distribution of the silver grains over some, but not all, of the striatal neurons as well as being concentrated around clusters of neurons. It appeared that the silver grains were absent from fibre tracks within the striatum. It appeared that PDE10A mRNA was not confined to regions close to the nucleus but was dispersed in cellular processes.

Huntingtin with an expanded polyglutamine tract (htt-HD) is expressed in neurons of the brain and body throughout development and during the lifetime of HD patients (The Huntington's Disease Research Collaborative, 1993; Ross, 1995). Transgenic HD mice express a portion of htt-HD and develop a phenotype with many of the symptoms of HD after a period of normal development and growth (Carter et al., 1999; Cha et al., 1998; Mangiarini et al., 1996). Using differential display RT PCR, northern blot and *in situ* hybridization, we have demonstrated that PDE10A mRNA levels decline in the striatum of HD mice. This specific member of the PDE multigene family is highly expressed in the striatum and olfactory tubercle of mice (Soderling et al., 1999) and rats (Fujishige et al., 1999) and in the caudate and putamen of humans (Fujishige et al., 1999; Loughney et al., 1999). The levels of PDE10A were the same in 5 week old wild-type and HD mice. PDE10A mRNA levels then began to decline and were almost undetectable in the striatum and olfactory tubercle by the time the mice reached 8 weeks of age. This time coincides with the onset of overt motor symptoms in the HD mice indicating that the loss of PDE10A in striatal neurons leads to

dysfunction of the nuclei that control movement. The R6/2 mice develop the HD phenotype in the absence of cell death. The decrease in PDE10A mRNA, therefore, is not due to the loss of PDE10A-expressing cells but rather a change in steady-state RNA levels that occurs due to the expression of mutant huntingtin.

The particular isoform that decreases in HD is PDE10A. PDE10A has been cloned from human lung and fetal brain cDNA libraries (Fujishige et al., 1999; Loughney et al., 1999). It appears that the presence of the expanded polyglutamine tract in huntingtin alters gene expression in the striatum, and that this is the mechanism by which only a small group of neurons in the striatum and cortex are rendered vulnerable to this ubiquitously expressed mutant protein.

EXAMPLE 8 - PDE10A is Highly Conserved Among Mammalian Species

The oligonucleotide (probe 1) complementary to the coding strand of the PDE10A transcript, was also used as an *in situ* hybridization probe against coronal brain sections derived from adult rats. FIG. 8 shows *in situ* hybridization analysis of adult rat brain sections using oligonucleotide probe 1 complementary to the coding-strand of PDE10A revealed that the pattern of expression of PDE10A is the same in rats and mice. The hybridization conditions used to detect the rat homologue of PDE10A in rat brain tissue differed from those used to detect the transcript in mice only in that the stringency of the post-hybridization washes were reduced.

No hybridization was observed in the rat striatum using the post-hybridization washes

employed following the *in situ* hybridization of mouse brain sections. However, when the stringency of the post-hybridization washes was lowered (2 x 60' in 1X SSC @ 42°C, 2 x 60' in 0.5X SSC @ 42°C, 2 x 60' in 0.25X SSC @ room temperature), the PDE10A oligonucleotide probe specifically labelled the adult rat striatum, nucleus accumbens and olfactory tubercule in a pattern indistinguishable from that observed in mouse brain sections. It appears, therefore, that a transcript which shares nucleotide sequence and expression pattern is present in both mice and rats. The evolutionary conservation of PDE10A suggests that it is important for normal function of the basal ganglia.

By northern blot, Fujishige et al. (1999) demonstrated that PDE10A is expressed in human fetal brain. The homology between mouse and human PDE10A is extremely high (data not shown).

EXAMPLE 9 - Analysis of PDE10A in Genomic DNA

Because the transgenic mice employed in this study have a copy of the human HD 5' UTR, exon 1 with expanded CAG repeat and 262 bp of the intron 1 that has been integrated into an undefined locus of the mouse genome, it was possible that the integration event disrupted the PDE10A gene preventing its expression in the HD mouse striatum. Genomic DNA was isolated from wild-type and HD mice and subjected to Southern blot analysis.

Genomic DNA was isolated from wild-type and HD mice and subjected to Southern blot analysis using pPDE10A as a hybridization probe. The size of the *Bam*HI and *Eco*RI fragments that are present in the transgenic R6/2 line that correspond to the insertion of the

human exon 1 gene fragment are 1.9 and 0.8 (*BamHI*) and 1.9 (*EcoRI*) kb. Analysis of the size of the fragments that hybridized with pPDE10A demonstrated that there was no difference in the size of the hybridizing fragments between the wild-type and HD mice. FIG. 9 shows the genomic DNA restriction fragments that hybridized with pPDE10A were the same in wild-type and HD mice. The size of the hybridizing *BamHI* and *EcoRI* fragments in each genomic DNA sample is approximately 8 kb and 3 kb, respectively. If the 1.9 kb *SacI-EcoRI* HD gene fragment integrated into the genome within the *BamHI* and *EcoRI* fragments that hybridized with the DHDM cDNA cloned insert, the sizes of the HD hybridizing bands would have been distinct from those of the wild-type. This Southern blot analysis indicates that the gene encoding PDE10A is present as a single-copy in the mouse genome. The numbers at the left of the blot are the relative mobility of molecular weight markers (1 kb ladder, BioRad).

The PDE10A cDNA has since been cloned using a bioinformatics search strategy involving screening of the expressed sequence tag (EST) database for novel PDE cDNA clones. Independently, the mouse PDE10A cDNA was identified after an EST search for novel PDEs with conserved cGMP binding domains (Soderling et al., 1999). The rat isoforms of PDE10A and splice variants have also been described (Fujishige et al., 1999). Human, mouse and rat PDE10A splice variants differ in their 5' untranslated and part of the 5' coding region but are identical in the coding region when the various splice variants are compared within each species. The human, mouse and rat PDE10A coding regions contain 779, 779 and 794 amino acids, respectively, encoding a protein of approximately 88.5 kDa.

EXAMPLE 10 - Distribution of PDE10A

In mouse, PDE10A mRNA was detected in testis and to a much lesser extent in brain but not in heart, spleen, lung, liver, skeletal muscle, kidney, ovary, pancreas, smooth muscle, eye or in total RNA isolated from 7, 11, 15 or 17 day old embryo (Soderling et al., 1999). This data agrees with the PDE10A mRNA pattern of distribution that we observed in wild-type and pre-symptomatic HD mice. In mice, two different size transcripts are detected in northern blots using the coding region as a probe. In mouse testis, the most abundant transcript is approximately 4 kb. A 9.5 kb transcript was also detected in mouse testis. It appears that the most abundant transcript in mouse brain is 9.5 k. Similarly, two sized PDE10A transcripts were observed in rats, however, it appears that, in rat, the 4 kb mRNA is expressed exclusively in testis while the 9.5 kb mRNA is expressed exclusively in brain (Fujishige et al., 1999). Within the brain, the rat PDE10A mRNA was expressed in striatum and olfactory tubercle and not cortex, cerebellum, hippocampus, midbrain or brainstem. In humans, PDE10A is expressed in the caudate, putamen and testis. As was observed in rodents, mRNAs of approximately 4 and 10 kb hybridized with the PDE10A probe. Again, it appears that, although both sized transcripts are present in brain and testis, the larger mRNA is predominant in the caudate and putamen and the smaller sized transcript is present in the testis. Each of the mouse, rat and human PDE10A sequences are not longer than 4 kb and span the coding region and parts of the 3' UTR. The difference in abundance of the short and long transcript in the testis and brain, respectively, in all three species suggest that the 3' UTR functions to provide transcript stability in the brain. As such, we present the complete sequence of the brain-specific transcript of PDE10A derived from mouse.

EXAMPLE 11 - Modulating Activity of PDE10A Using cGMP-PDE Activity

Cyclic nucleotides are the predominant second messengers that activate cellular signaling pathways (Beavo, 1995; Conti and Jin, 1999). The concentration of intracellular cyclic nucleotides is dependent on their rate of synthesis by adenyl and guanyl synthase, the rate of efflux from the cell, and the rate of degradation. PDEs hydrolyze cAMP and cGMP limiting both the duration and amplitude of the cyclic nucleotide signal (Beavo, 1995; Conti and Jin, 1999). In mammals, PDEs are encoded by a large multigene family. The various PDE family members have tissue-specific patterns of expression (Conti and Jin, 1999). PDEs have also been described in Caenorhabditis, Drosophila, Dictyostelium, Saccharomyces, Candida and Vibrio species demonstrating that this enzyme has been conserved throughout evolution. In mammals, PDEs are encoded by at least 10 gene families, each composed of one or more genes. In addition, numerous splice variants of individual gene family members have been described. These splice variants alter the 5' domain of the protein but share identical nucleotide binding and catalytic domains. The catalytic domain, found in the carboxyterminus of the enzyme, is ~ 275 amino acids and highly conserved in amino acid sequence in all PDEs. In total, it appears that there are ~50 PDEs expressed within the mammalian body. Some PDEs are expressed in multiple tissues while others have a very limited tissue-specific distribution (Conti and Jin, 1999).

PDE gene families differ with respect to their affinity for cAMP and cGMP and their dependence on calcium and calmodulin (Beavo, 1995). Moreover, some PDEs are inhibited or activated by binding cyclic nucleotides to a non-hydrolytic site. For example, PDE2A has a lower K_m for cGMP than cAMP although it hydrolysed both nucleotides. The binding of

cGMP to an allosteric activator site within PDE2 enhances the rate of catalysis of cAMP. PDE2 is, therefore, a cGMP-stimulated cGMP and cAMP phosphodiesterase (Beavo, 1995). Conversely, the affinity of PDE4 for cAMP is much greater than for cGMP and PDE4 activity is not affected by cGMP or calmodulin (Beavo, 1995). The differences in substrate preference, modulation of activity and tissue-specific patterns of expression suggest that subtle alterations in the relative levels of cAMP and cGMP mediated through the action of various PDEs lead to a wide range of responses to extracellular signals.

cGMP-PDE activity of compounds is measured using a one-step assay adapted from Wells at al. (Wells, J. N., Baird, C. E., Wu, Y. J. and Hardman, J. G., *Biochim. Biophys. Acta* 384:430 (1975)) and adopted by Beavo et al, U.S. Patent No. 6,037,119. The reaction medium contains 50 mM Tris-HCl, pH 7.5, 5 mM Mg-acetate, 250 ug/ml 5'-Nucleotidase, 1 mM EGTA and 0.15 uM 8-[H³]-cGMP. The enzyme used is a human recombinant PDE V (ICOS, Seattle U.S.A.).

Compounds of interest are dissolved in DMSO finally present at 2% in the assay. The incubation time was 30 minutes during which the total substrate conversion did not exceed 30%.

The IC ₅₀ values for the compounds examined are determined from concentration-response curves using typically concentrations ranging from 10 nM to 10 uM. Tests against other PDE enzymes using standard methodology also show compounds highly selective for the cGMP specific PDE enzyme.

Rat aortic smooth muscle cells (RSMC) are prepared according to Chamley et al. in *Cell Tissue Res.* 177:503-522 (1977) and used between the 10th and 25th passage at confluence in 24-well culture dishes. Culture media is aspirated and replaced with PBS (0.5 ml) containing the compound tested at the appropriate concentration. After 30 minutes at 37° C, particulates guanylate cyclase are stimulated by addition of ANF (100 nM) for 10 minutes. At the end of incubation, the medium is withdrawn and two extractions were performed by addition of 65% ethanol (0.25 ml). The two ethanolic extracts are pooled and evaporated until dryness, using a Speed-vat system. c-GMP was measured after acetylation by scintillation proximity immunoassay (AMERSHAM). The EC₅₀ values are expressed as the dose giving half of the stimulation at saturating concentrations.

EXAMPLE 12 - Selected Modulators of PDE10A Activity

The catalytic domain of PDE10A is most similar in amino acid sequence to PDE5A, PDE2A, PDE6B and PDE6A. These members of the PDE family each contain a cGMP binding sequence that is not observed in other PDE family members. The non-catalytic cGMP binding sites (GAF) domains found in PDE2, 5 and 6 are also found in PDE10. At least for PDE2, this site acts as an allosteric activator for cAMP hydrolytic activity. The GAF domain of PDE10A binds other small molecules that act as allosteric activators. PDE10A is a cAMP and cAMP-inhibited cGMP PDE (Fujishige et al., 1999; Fujishige et al., 1999; Loughney et al., 1999; Soderling et al., 1999).

Attenuation of the production of cAMP, may ameliorate the symptoms of HD and positively affect gene expression. Pharmaceutically acceptable modulators of cAMP include quinpirole,

alloxan, miconazole nitrate, MDL-12330A, and tetracyline derivatives such as demeclocycline and minocycline.

Compounds which are potent and selective modulators of cGMP-specific PDE, and are useful in a variety of therapeutic areas are taught by Daugan et al, U.S. patent No. 5,981,527, PCT publication No. WO 00/15639 to Icos Corporation and PCT publication No. WO 00/15228 to Icos Corporation, which are incorporated herein by reference. Such compounds include, for example:

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-methyl-pyrazino[2', 1':6,1]pyrido[3,4-b]indole-1,4-dione,

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-pyrazino[2',1':6,1]py rido[3,4-lindole-1,4-dione,

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-isopropyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione,

(3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-3-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione, and

(3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2,3-dimethyl-pyraz ino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione.

PDE1B1 is expressed throughout the brain and is most abundant in the striatum, nucleus accumbens and olfactory tubercle (Polli and Kincaid, 1994; Yan et al., 1994). PDE1B is a cGMP, Ca/calmodulin-dependent PDE. Therefore, PDE1B and 10A are both expressed in the majority, but not all, striatal neurons and, it is likely that both genes are co-expressed in a subset of striatal projection neurons. Selective inhibitors for PDE1 include KS-505, IC224,

and SCH 51866. Of these inhibitors, it appears that SCH 51866 has a ten-fold higher Km for PDE1 than PDE10 (Soderling et al., 1999). The non-specific PDE inhibitor IBMX is a potent inhibitor of PDE10A. Dipyridamole and SCH51866 had the highest potency of inhibitors tested on PDE10A activity. Dipyridamole was considered to be a PDE5- and PDE6-specific inhibitor, however, the Km for dipyridamole is 10 times higher for PDE10A than the other PDEs (Soderling et al., 1999). Selective inhibitors of PDE5, 2, 3 and 4 had much greater IC50 for PDE10 (Soderling et al., 1999).

EXAMPLE 13 - Clinical use of PDE10A Modulator

A 38 year-old female was admitted to hospital from a long-term care facility due to progressive deterioration of her physical and mental symptoms caused by Huntington's disease. The patient had been diagnosed with Huntington's disease at age 26. Prior to admission to the hospital, she had become increasingly aggressive and uncooperative.

Moreover, there appeared to be an increase in the number of psychotic episodes. SPECT showed no abnormality of brain blood flow but MRI showed bilateral caudate atrophy as well as global atrophy of the cerebrum and corpus callosum.

The patient had been stable for a number of years on the antipsycotic haloperidol (3 mg/day). For the last two years, the haloperidol had been replaced by olanzapine (2.5-7.5 mg/day).

Minocycline, a tetracycline derivative, was administered at 50 mg twice daily for 7 days, followed by 100 mg twice daily for 7 days and finally 200 mg twice daily for 5 weeks. After 5 weeks of 200 mg twice daily minocycline administration, there was a mild improvement

compared to the baseline clinical global assessment made at the time of admission. The minocycline treatment was suspended for 7 days. Due to a significant increase in the number of aggressive incidence and decrease in cooperativity, minocycline (200 mg twice daily) treatment was resumed. The patient responded within 3 days to the resumed minocycline-treatment with a return to mild-improvement compared to the baseline clinical global assessment made at the time of admission. Minocycline (200 mg twice daily) treatment will continue indefinitely. The improvement in behaviour and decrease in apparent psychosis has allowed for the transfer of the patient from the acute care facility back to long-term care.

While the present invention has been described in terms of specific embodiments, it is understood that variations and modifications will occur to those skilled in the art.

Accordingly, only such limitations as appear in the appended claims should be placed on the invention.

- Non-patent Literature Cited
- Atschul, S. F. et al.(1990), J. Molec. Biol. 215: 403.
- Babity, J.M., Armstrong, J.N., Plumier, J.-C., Currie, R.W., and Robertson, H.A. (1997a) A novel seizure-induced synaptotagmin gene identified by differential display. *Proc. Nat. Acad. Sci. U.S.A.* 94: 2638-2641.
- Babity, J.M., Newton, R.N., Guido, M.E. and Robertson, H.A. (1997b) The Application Of Differential Display To The Brain: Adaptations For The Study Of Heterogeneous Tissue.

 Methods Mol Biol, 85 (1997) 285-95.
- Barney (1991) in PCR Methods and Applications, Aug., Vol. 1(1), page 5.
- Beavo, J. (1995). Cyclic nucleotide phosphodiesterase: functional implications of multiple isoforms. Physiol. Rev. 75, 725-748
- Beretta, S., Robertson, H. and Graybiel, A. (1992) Dopamine and glutamate agonists stimulate neuron-specific expression of Fos-like protein in the striatum. *J. Neurophysiol.* 68, 767-777
- Bennett, D. et al., (1995) Journal of Molecular Recognition, Vol. 8 52-58.
- Berke, J.D., Paletzki, R.F., Aronson, G.J., Hyman, S.E. and Gerfen, C.R. (1998) A Complex Program of Striatal Gene Expression Induced by Dopaminergic Stimulation. *J. Neurosci.* 18: 5301-5310.
- Caine E.D., Hunt R.D., Weingartner H., Ebert M.H. (1978) Huntington's dementia. Clinical and neuropsychological features. *Arch-Gen-Psychiatry*. 35, 377-84.
- Carillo, H., and Lipman, D.(1988), SIAM J. Applied Math., 48: 1073.
- Carter, R. J., Lione, L. A., Humby, T., Mangiarini, L., Mahal, A., Bates, G. P., Dunnet, S. B., and Morton, A. J. (Apr. 1999). Characterization of progressive motor deficits in mice transgenic for the human Huntington's diease mutation. J Neurosci 19, 3248-57

Cha, J.-H. J., Kosinski, C.M., Kerner, J.A., Alsdorf, S.A., Mangiarini, L., Davies, S.W., Penny, John B., Bates, G.P., Young, A.B. (1998) Altered brain neurotransmitter receptors in transgenic mice expressing a portion of an abnormal human Huntington disease gene. *Proc. Natl. Acad. Sci. USA* 95: 6480-6485.

- Conolly, B. A. (1987) Nucleic Acids Res. 15:15(7): 3131.
- Conti, M., and Jin, S. L. (1999). The molecular biology of cyclic nucleotide phosphodiesterases. Prog Nucleic Acid Res Mol Biol 63, 1-38
- Crino, P., Khodakhah, K., Becker, K., Ginsberg, S., Hemby, S. and Eberwine, J. (1998)
 Presence and phosphorylation of transcription factors in developing dendrites. *Proc. Natl. Acad. Sci. USA* 95: 2313-2318
- Davies, S.W., Turmaine, M., Cozens, B.A., DiFiglia, M., Sharp, A.H., Ross, C.A., Scherzinger, E., Wanker, E.E., Mangiarini, L. and Bates, G.P. (1997) Formation of neuronal intranuclear inclusions underlies the neurological dysfunction in mice transgenic for the HD mutation. *Cell* 90, 537-548.
- Davis et al., (1986) Basic Methods in Molecular Biology.
- Denovan-Wright, E.M., Newton, R.A., Armstrong, J.N., Babity, J.M., Robertson, H.A. (1998) Acute administration of cocaine, but not amphetamine, increases the level of synaptotagmin IV mRNA in the dorsal striatum of rat. *Mol. Brain Res.* 55, 350-354.
- Denovan-Wright, E.M., Gilby, K.L., Howlett, S.E. and Robertson, H.A. (1999) Cloning of differentially expressed brain cDNA. In *Differential Display A Practical Approach*,
 H.A. Robertson and R.A. Leslie, Eds. Oxford University Press (in press).
- Devereux, J., et al.(1984), Nucleic Acids Research 12(1): 387.
- Douglass, J., McKinzie, A.A. and Couceyro, P. (1995) PCR differential display identifies a rat brain mRNA that is transcriptionally regulated by cocaine and amphetamine. *J.*

- Neurosci. 15, 2471-2481.
- Douglas, J. and Daoud, S. (1996) Characterization of the human cDNA and genomic DNA encoding CART: a cocaine- and amphetamine-regulated transcript. *Gene* 169, 241-245
- Eberwine, J. (1996) Amplification of mRNA populations using aRNA generated from immobilized oligo(dt)-T7 primed cDNA. *Biotechniques* 20: 584-589
- Eberwine, J., Yeh, H., Miyashiro, K., Cao, Y., Nair, S., Finnell, R., Zettel, M., Coleman, P., (1992) Analysis of gene expression in single live neurons. *Proc. Natl. Acad. Sci. USA* 89: 3010-3014
- Fujishige, K., Kotera, J., Michibata, H., Yuasa, K., Takebayashi, S., Okumura, K., and Omori, K. (June 25, 1999). Cloning and characterization of a novel human phosphodiesterase that hydrolyzes both cAMP and cGMP (PDE10A). J Biol Chem 274, 18438-45
- Glass, M., Faull, R.L.M. and Dragunow, M. (1993) Loss of cannabinoid receptors in the substantia nigra in Huntington's disease. *Neuroscience* 56: 523-527
- Good et al (1977), Nucl. Acid Res 4:2157.
- Gribskov, M. and Devereux, J., eds., Sequence Analysis Primer, M Stockton Press, New York, 1991.
- Griffin, A. M., and Griffin, H. G., eds., Computer Analysis of Sequence Data, Part I, Humana Press, New Jersey, 1994.
- Group THDCR (1993) A novel gene containing a trinucleotide repeat that is extended and unstable on Huntington's disease chromosomes. *Cell* 72: 971-983.
- Hardy, J. and Gwinn-Hardy, K. (1998) Genetic classification of primary neurodegeneraive disease. *Science* 282: 1075-1079
- Hinnen et al., (1978) PNAS USA 75:1929-1933.

Houbenweyl, (1987) *Methods of Organic Chemistry*, ed. E. Wansch, Vol. 15 I and II, Thieme, Stuttgart.

- Innis, M. A. and D. H. Gelfand (1989), PCR Protocols, A guide to Methods and Applications, M. A. Innis, D. H. Gelfand, J. J. Shinsky and T. J. White eds, pp 3-12, Academic Press.
- Johanson, K. et al.(1995), The Journal of Biological Chemistry, Vol. 270, No. 16, pp 9459-9471.
- Klement, I.A., Skinner, P.J., Kaytor, M.D., Yi, H., Hersch, S.M., Clark, H.B., Zoghbi, H.Y. and Orr, H.T. (1998) Ataxin-1 nuclear localization and aggregation: role in polyglutamine-induced disease in *SCA1* transgenic mice. *Cell* 95: 41-53.
- Lesk, A. M., ed., *Computational Molecular Biology*, Oxford University Press, New York, 1988.
- Liang, P. and Pardee, A.B. (1992) Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257: 967-971.
- Livesey, F.J., O'Brien, J.A., Li, M., Smith, A.G., Murphy, L.J. and Hunt, S.P. (1997) A

 Schwann cell mitogen accompanying regeneration of motor neurons. *Nature* 390: 614-618.
- Livesey, F.J. and Hunt, S.P. (1996) Identifying changes in gene expression in the nervous system: mRNA differential display. *Trends Neurosci.* 19: 84-88
- Loughney, K., Snyder, P. B., Uher, L., Rosman, G. J., Ferguson, K., and Florio, V. A. (June 1999). Isolation and characterization of PDE10A, a novel human 3', 5'-cyclic nucleotide phosphodiesterase. Gene 234, 109-17
- Ludlow, C.L., Connor, N.P., Bassich, C.J. (1987) Speech timing in Parkinson's and Huntington's disease. *Brain-Lang*, 32, 195-214.
- Mangiarini, L., Sathasivam, K., Seller, M., Cozens, B., Harper, A., Hetherington, C. Lawton,

M., Trottier, Y., Lehrach, H., Davies, S. W. and Bates, G. P. (1996) Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice. *Cell* 87: 493-506.

- Merrifield, (1964), J. Am. Chem. Assoc. 85:2149-2154.
- Okano, J., (1988) Neurochem. 56: 560.
- Pardee, A.B. (1997) Complete genome expression monitoring: the human race. *Nat. Biotechnol.* 15: 1343-1344
- Polli, J.W., and Kincaid, R. L. (1994). Expression of a calmodulin-dependent phosphodiesterase isoform (PDE1B1) correlates with brain regions having extensive dopaminergic innervation. J. Neurosci. 14, 1251-1261
- Qu,-D., Ludwig,-D.S., Gammeltoft, S., Piper, M., Pelleymounter, M.A., Cullen, M.J. Mathes, W.F., Przypek, R., Kanarek, R. and Maratos-Flier, E. (1996) A role for melanin-concentrating hormone in the central regulation of feeding behaviour. *Nature* 380: 243-247.
- Richfield, E.K. and Herkenham, M. (1994) Selective vulnerability in Huntington's disease: preferenctial loss of cannabinoid receptors in lateral globus pallidus. *Ann. Neurol.* 36: 577-584.
- Ross, C.A., (1997) Intranuclear neuronal inclusions: a common pathogenic mechanism for glutamine-repeat neurodegenerative diseases? *Neuron* 19: 1147-1150.
- Saiki et al., (1986) Nature, 324: 163-166.
- Sambrook, et al, (1989), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Saudou, F., Finkbeiner, S., Devys, D. and Greenberg, M.E. (1998) Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear

- inclusions. Cell 95: 55-66.
- Sisodia, S.S., (1998) Nuclear inclusions in glutamine repeat disorders: are they pernicious, coincidental or beneficial? *Cell* 95: 1-4.
- Smith, D. W., ed., *Biocomputing: Informatics and Genome Projects*, Academic Press, New York, 1993.
- Soderling, S. H., Bayuga, S. J., and Beavo. J. A. (June 1999). Isolation and characterization of a dual-substrate phosphodiesterase gene family: PDE10A. Proc Natl Acad Sci U S A 96, 7071-6
- von Heinje, G., Sequence Analysis in Molecular Biology, Academic Press, 1987.
- Yan, C., Bentley, J. K. Sonnenburg, W. K., and Beavo, J. A. (1994). Differential expression of the 61 kDa a 63 kDa calmodulin-dependent phosphodiesterase in the mouse brain. J. Neurosci. 14, 973-984
- Young, A.B., Penney, J.B., Starosta-Rubinstein, S., Markel, D.S., Berent, S., Giordani,
 B., Ehrenkaufer, R., Jewett, D., Hichwa, R. (1986) PET scan investigations of
 Huntington's disease: cerebral metabolic correlates of neurological features and
 functional decline. *Ann-Neurol*. 20, 296-303.

We claim:

1. A composition for treating a CAG repeat disorder comprising a compound which modulates PDE10A expression and a pharmaceutically acceptable carrier.

- 2. A composition as claimed in claim 1, wherein said compound is selected from the group consisting of: quinpirole, alloxan, miconazole nitrate MDL-12330A, and tetracyline derivatives such as demeclocycline.
- 3. A composition as claimed in claim 1, wherein said compound is selected from the group consisting of:

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-methyl-pyrazino[2', 1':6,1]pyrido[3,4-b]indole-1,4-dione,

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-pyrazino[2',1':6,1]py rido[3,4-lindole-1,4-dione,

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-isopropyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione,

(3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-3-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione,

(3S,6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2,3-dimethyl-pyraz ino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione.

4. A composition as claimed in claim 1, wherein said compound is selected from the group consisting of: KS-505, IC224,SCH 51866, IBMX and Dipyridamole.

5. A composition as claimed in any one of claims 1 to 4, wherein said disorder is Huntington's disease.

- 6. The use of a composition as claimed in any one of claims 1 to 5 for treating a CAG repeat disorder comprising administering said composition to a subject in need of such treatment.
- 7. The use of a composition of claim 6 for treating Huntington's disease comprising administering said composition to a subject in need of such treatment.
- 8. A method for identifying a compound which inhibits or promotes a CAG repeat disorder, comprising the steps of:
- (a) selecting a control animal having PDE10A and a test animal having PDE10A;
- (b) treating said test animal using a compound; and,
- (c) determining the relative quantity of RNA corresponding to PDE10A, as between said animals.
- 9. A method of claim 8, wherein said animal is a mammal.
- 10. A method of claim 9, wherein said mammal is a mouse.
- 11. A method of claim 10, wherein said mouse is R6/2 transgenic mouse.
- 12. A method of any one of claims 8 to 11, wherein said CAG repeat disorder is Huntington's disease.

13. A method for identifying a compound which inhibits or promotes a CAG repeat disorder, comprising the steps of:

- (a) selecting a host cell containing PDE10A;
- (b) cloning said host cell and separating said clones into a test group and a control group;
- (c) treating said test group using a compound; and
- (c) determining the relative quantity of RNA corresponding to PDE10A, as between said test group and said control group.
- 14. A method of claim 13, wherein said CAG repeat disorder is Huntington's disease.
- 15. A method for detecting the presence of or the predisposition for a CAG repeat disorder, said method comprising determining the level of expression of RNA corresponding to PDE10A in an individual relative to a predetermined control level of expression, wherein a decreased expression of said RNA as compared to said control is indicative of a CAG repeat disorder.
- 16. A method of claim 15, wherein said CAG repeat disorder is Huntington's disease.
- 17. A method of claim 15 or 16, wherein said expression is measured by in situ hybridization.
- 18. A method of claim 15 or 16, wherein said expression is measured using a polymerase chain reaction.

19. A method of claim 15 or 16, wherein said expression is measured using a DNA fingerprinting technique.

HD WT

123456



Figure 1

Striatum Cortex

HD WT HD WT

123456123456

9.5

7.5




4.4

2.4

Figure 2

Figure 3

5 '		11	21	31	41
-	TGTATGGGAA ACATACCCTT	TAGTGTTTCC	CATATGATCTO	TTGTCTGGAG	TATATGCTAC
ز	ACATACCCTT	ATCACAAAGC	TATACTAGAC	CAACAGACCTC	CATATACGATG
				81	probe 1
5 '		61	71	81	91
C 1	ATGTTCATTT. TACAAGIAAA	ACTGTACAAA	AACCCAGTGC	CAGCTGATGAT	GCAAAGCAGT
51	TACAAGTAAA	TGACATGTTT	TTGGGTCAC	TCGACTACTA	ACGTTTCGTCA
5 '		11	21	31	41
101	CTCTCTCTGT GAGAGAGACA	GTACAGTGCC	CCACCTATTI	TAAAAATCACG	TACTTGCCCA
101	GAGAGAGACA	CATGTCACGG	GGTGGATAAA	ATTTTTAGTGC	ATGAACGGGT
5 '	GAACACTGTGA CTTGTGACAC	51	71	81	91
151	GAACACTGTG	AAACACTTAA	CATAAGAACA	AACGCAGCGT	CIGGATICII
101	CTTGTGACAC'	TTTGTGAATT	'GTATTCTTGT	TIGCGICGCA	GACCTAAGAA
				21	41
5 '	TCCAAGGAGAC AGGTTCCTCTC	11 proce 2	21	31	41
201	TCCAAGGAGA	GCAGCTTTCT	CCACAGGAAC	ACAGTAACAA	AAGAGGTCCG
2.01	AGGTTCCTCT	CG'I'CGAAAGA	GGTGTCCTTG	TGTCATTGTT	TICICCAGGC
			•7 1	81	91
5 '	,	51	, T	CYCCCCV TIVC	
251	CCGCCATCCA C	ACCCAGCCA	AGACACCTCA	CARCACCATAG	CCTCTTCCACC
	GGCGGTAGGT	alegeleet.	TCIGIGGAGI	CICCGGIAIC	CCIGIIGAG
- 1	1	1	21	31	41
5 '		. T . T	aayaayaaa T	.CXCXGGTCCC	
301	CTTGCTGGCCA GAACGACCGGT	THCHCCIGCI	CCTCCTCCCC	CTCTCCAGGG	TCGTTGACTA
	GAACGACCGG 1	. I G I G GACGA	CCICGICCC	٠,٠٠٠ مينان د	
5 '	6	1	71	81	91
3	CCTCAGTGGAI GGAGTCACCTA	יב יכככידרידכר מ	GCCAAAGCCT	TAATGGGCTC	TCTTTTGAAG
351	CCICAGIGGAI		CGGTTTCGGA	ATTACCCGAG	AGAAAACTTC
5 '	1 GGGAAAGAAAG CCCTTTCTTTC	1 :	21	31	41
_	CCCDAAGAAAG	AATTTCAAG	CTTATGATAT	CCAATATTAT	TATAGTTGAT
401	CCCTTTCTTTC	TTAAAGTTC	GAATACTATA	GGTTATAATA	ATATCAACTA
5 '	6		71	81	91
_	GAGTTAGTAAA	TTCCAAAAA	AAAAA		
451	GAGTTAGTAAA CTCAATCATTT	AAGGTTTTŢ	PTTTT		



WT HD

Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

Figure 10


5	ſ	11	21	31	41
_	CACTGAAGO	CTGGTCCACGT	CTATAAAC	AGG TGACACTGO	ectgcagcaaa <i>i</i>
	1 GTGACTTC	EA CCAGGTGC	GATATTTG	TCCACTGTGAC	CGACGTCGTTT
				0.7	
5		61	71		91
5	AGCCATTCG	SA TCCACACAA TT NGCTCTCTT	MATTGATCT TTNNCTNCN	CTATCATCTTC AGATAGTAGAAC	CAATCIGAATI
_	- TCGGTAAGC	TAGGIGIGII	IAACIAGAA	ANDATONI NOP	CITAGACITA
5	t	11	21		41
	CCACCCACC	AGCAGTATGT	AAGACGAC	CGTTTAATTCAG	GCATTCCGAAG
10	1 CGTCCCTCC	TCGTCATACA	TTCTGCTGG	CAAATTAAGTO	CGTAAGGCTTC
	·	~ .	71	81	91
5	· · · · · · · · · · · · · · · · · · ·	61	71	אר א א ש א דוט אר א א א ש א דוט א א א א פר	
15	1 GCATGAGCG	CATGGATTCT	G TCACCAAG C NGTGGTTC	CGTATAAAAGG CGCATATTTTCC	TCCCIGGCAII TCCCACCCTAA
	- CGTACTCGC	G IACCIAAGA	CAGIGGIIC	GCAIAIIIICC	I OUCHCCGIAN
5 '		11	21	31	41
	<u> </u>	ል ጥር <mark>ል</mark> ርርርር <mark>ል</mark> ርጥ	GTTTTTGCT	GTAGAAGTAGG	GATTTTACAGA
20	1 CCCTTTGGA	TACTGCCTGA	CAAAAACGA	CATCTTCATCC	CTAAAATGTCT
				0.7	0.1
5 '		61 ~~~~~~	71	81	91
25	AGTCTCCTT	GAATTTGCCC	r GCCTGGGG	CAGTTTTGCAG GT CAAAACGTC	T CCTTGC A CCC
	TCAGAGGAA	CITAAACGGG	ACGGACCCC	GI CAMMACGIC	1 CC11 COACGO
5 '		11	21	31	41
	AGAGATTTA:	rTGGCTGGTC	GTCTCTTG	TGAAATAGTAT	CATGTGAGAAA
30:	TCTCTAAAT?	AACCGACCAGT	CAGAGAAC	ACTTTATCATA	GTACACTCTTT
		~ -	71	81	91
5 '		61 ************************************	\ T	AAGACCTTTGC	א א האתהמתההה א א האתהמתהמה
351	CAGTTTGTAC	₹₩₩₩₩₩₩Ċ ₽₩₽₩₽₽₽₽₽₽	TACCIGGG	TTCTGGAAACG'	TTGTAACAAGG
	GICAAACAIC	JIIIIIIOMI	HI CONCO	110100111100	
5 '		11	21	31	41
407	TTCCATGGGC	CAAGACTCAG	TTAGGAGG	CATAAATCTGC	CCGGAATAAAC
401	AAGGTACCCG	GTTCTGAGTC	AATCCTCC	GTATTTAGACG(GCCTTATTTG
		C1	71	81	91
5'	ma aaaaaaaaaa	61	፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞፞ኯኯኯኯኯ ፞ኯኯኯኯኯኯኯኯ	AT AATTTGGTTT	TAGAATTCAC
451	TAGGCCAGGA TAGGCCAGGA	ATGTCGGTAC	AAATCAATT	TATTAAACCAA!	ATCTTAAGTG
	Arccourcer	111010001110			•
51		11	21	31	41
E 0.1	ACAGGCAGGA	TTGGTTTTTT	TGTGTCTTG	31 GCAAGTGGAGC CGTTCACCTC	CATATTTAACA
201	TGTCCGTCCT	AACCAAAAAA	ACACAGAAC	CCGTTCACCTCC	TATAAATTGT
5 '	*	61	71	81	91
5'	תה מה מממה היה	CCDDTCCTCC	᠘ᡙᢕᡎᡎᢧᢙᢕᡎ ᠘ᠴ	TTTCCCACCCI	
551	ATGTCCGTAC	CCTTAGGACG	GAGAATCGA	TTTCCCACCCI AAAGGGTGGGA	GAACAGAGTG
	111010001110				
5'			21	31	41
601	CAAGTTTTTT	CTCTCCAAAG	GTTTCCAGG	A ATTTCTCATT TTAAAGAGTAA	AATGGCTGAT
007	ርጥጥሮልልልልልል	FAGAGGTTTC	CAAAGGTCC	TTAAAGAGTAA	TTACCGACTA

5	t	61	71	81	91
-	GCAAACTTA	GTGAATAATA	ATGAATATAA	A CAATGCTCA	CCTCACCAAAA
65	1 CGTTTGAAT	CACTTATTAT	TACTTATATT	TGTTACGAGT	CCTCACCAAAA GGAGTGGTTTT
5 '	ı	11	21	31	41
70	TTATATTAT	TTGCAGTCAT	TTGTGATAAC	A CAAATTTTA'	TCGCAATGGTT
70	ATATAATAA ¹	AACGTCAGTA	AACACTATTG	TGTTTAAAAT	AGCGTTACCAA
		6 7	71	81 .	91
5 '		61	. / T マッマックのころ	ᠬᠳᢧᡎᢕᡎᡎᡎᡎᢗᡸ	TGTGGTTGTT
75	ATTATTTAA:	TTTGTGGCCA	CACACIGIGG	A ATAGAAAAC	AACACCAACAA
	- TAATAAATI				
5 '		11	21	31	41
5	መርም <u>ሮ</u> አር አ አ አ አ	^ \ TGTTCTTGG	A TATGTAAGT	GCCAATACCA	STGTGAAGTAT CACACTTCATA
80		racaagaacc'	TATACATTCA	CGGTTATGGT	CACACTTCATA
	AONCICILI.				
5 '		61	71	81 .	91
		CAGCAAAAT	A CAGCCTAAG	GTTTGTAAAC	ATCAATTCTAT
85.	ACTAGGGCCC	CGTCGTTTTA!	rgtcggattc	CAAACATTTG:	ragttaagata
•			21		41
5 '		11	21 - ~- ~- ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	31	
903	CTCAGTTCAT	CAGAGGGCC	CHAGAAGCTG	CCCCCTCACIG	TAAAGTAAAGT ATTTCATTTCA
, ,	GAGTCAAGTA	AGTCTCCCGGY	CTCTTCGACC	3CCCCG1 CAC	11110111101
		61	71	81	91
5 '	* maamaaaaa	፞ፚፚኯፚፚኯፚዼኯ ፞	* A GCCTCCCG	CTGAAGAGTG	ACCAGTGCTG
951	TATGCIGGGCI	CCACCACCAC	TCGGAGGGC	GACTTCTCAC	FACCAGTGCTG CTGGTCACGAC
	IACGACCCCI				•
5.1		11	21	31	41
	GCCCGACGGA	TCGCTGAGAI	ATTCTCCCAT	'AATGGCAAA <i>I</i>	AAATAGGCAG
1001	CGGGCTGCCT	AGCGACTCTA	TAAGAGGGT	ATTACCG1"1"1"1	AAATAGGCAG TTTTATCCGTC
				•	91
5 '		61	71	℧ℒ ℸℼⅆⅆⅆⅆⅆⅆⅆⅆ	CATGTGTTAGC
1051	TTTGATGTGA	CCTGTTTAGT	GIGGCICICC	ZACADAACTCC	TACACAATCG
1001	AAACTACACT	GGACAAAICA		MOMMMICIOC	,
5 '		11	21	31	41
5	a manana a man	.	CAGTGAACTC	TGCTCTTCCA	AGTGTGTTCA
1101	ATTTTTATT AAAAAATAAA.	ATATGAGTAG	GTCACTTGAG	ACGAGAAGGI	TCACACAAGT
			·	•	
5 '		61	71	81	91
7757	mama mamaam	AGATATATTA	GCACAGCCTG	CCTTCTGCTG	91 CACAACGCCT GTGTTGCGGA
TTOT	ACATACACGA'	TCTATATAAT	CGTGTCGGAC	GGAAGACGAC	GTGTTGCGGA
		4 4	21	31	41
5'		11	ᆇᆂ ᄺᅒᄺᄱᄳᄳᅒᄺᄺᅋ		TTCTGCTCTC
1201	TAGAGACCCG	CCTTTCAAT	CTCCD DTCCD	ACACGAGACA	AAGACGAGAG
	ATCTC1GGGC	CGGWWWGIIN	CI COMMI COM		
5 '	é	51	71	81	91
_	ጥጥ አርረጥርጥ አ አ ፤	CTATGGTGT	CAGTTTTAAT	AGAACAAAAG	TATGCATCTT
1251	AATCCAGATT	GATACCACA (GTCAAAATTA	TCTTGTTTTC	ATACGTAGAA
				•	

5 '	7 7	0.7	2.4	
5 ·	ርጥጥርርርርምምርያ ጉጉ	արդություն ՀՀ	בי. אינים אינים ביניים אינים אינים אינים בינים אינים בינים אינים בינים אינים אינים אינים אינים אינים אינים אינים א	41
1301 CG	GAACCGAACTCGGA	AAAGCAAAA	GTTACGACTGA	41 TCTCCCCTTTCTCT AGAGGGGAAAGAGA
5'	61	71	81	91
1351 CC7	rgtgctcaccttac Acacgagtggaatg	CTTT CCAGA GAAA GGTCT	GTGTA AGGGAC CACAT TCCCTG	AACTTTTAAGGAGG TTGAAAATTCCTCC
5 '	11	21	31	41
$1401 \frac{\text{CGT}}{\text{GCA}}$	GTCCCTGGTAGGG CAGGGACCATCCC	GCAT CCCTG CGTA GGGAC	TTCAC CAGGTG AAGTG GTCCAC	CCTGTCATCACCCC GGACAGTAGTGGGG
5 '	61		81	91
1451 ACT TGA	TGACTGACATCTA ACTGACTGTAGAT	CCTGGTGA(GGGACCACT(CTATGGGTTCC: SATACCCAAGG	PCTT GTTTGTAGGG AGAA CAAACATCCC
5 '	11	21	31	41
	•			TTCTGGTTCCCGGC AAGACCAAGGGCCG
5'	61	71	81	91
1551 ACG	61 CTTTGGTTTTGAAA JAAACCAAAACTTT	GTCTCTTCT CAGAGAAGA	CTGTATATTCC GACATATAAGG	TACCCTGCATTTG ATGGGACGTAAAC
5 1	11	21	31	41
1601 CTTT	TGTGTGGTGCTGAT ACACACCACGACTA	GCTGTGCGC CGACACGCG	AGTA GGATTCT TCAT CCTAAGA	TGGATGACTCTCC ACCTACTGAGAGG
5 1	61	71	81	91
1651 ATCA	61 GTCACAGACTCCC CAGTGTCTGAGGG	CCTGTTGCA GGA CAACGT	AAGTGTCAGGC FTCA CAGTCCG	TGA CTCGACAGTC ACTGAGCTGTCAG
5 '	11	21	31	41
1701 ACCG TGGC	TAAAAT CTGAGTCA ATTTTA GACTCAG	AGT CACACAC CCAGTGTGTC	CAGG CTGTCAG STCCGACAGTC	CCA CGGCTTCCAC GGTGCCGAAGGTG
5'	61	71.	81	91
	61 ATGGCTATTCTATT FACCGATAAGATA			
5 '	11	21	31	41
1801 ACTGO	11 SCATTA TCTATGCT CGTAAT AGATACGA	AAGTTGAAA TTCAACTTT	TCAGGAGTGCC AGTCCTCACGG	CCAGCAGAGCCCA GGTCGTCTCGGGT
5'	61	71	81	91
1851 TCATT	CTCACTGTCTTTG GAGTGACAGAAAC	AAACAAAGC' TTTGTTTCG	TGTACGGTTTG ACATGCCAAAC	ATCGATGAACGT TAGCTACTTGCA
5'	11	21	31	41
1901 ATTTA TAAAT	AAGCATTTCATGC: TTCGTAAAGTACG:	AATGACAAA(FTACTGTTT(ETGCTCAGTAG CACGAGTCATC	TGGAAGGCAGGC ACCTTCCGTCCG


. 5	61	71	81	91	
_	ምርምር እ <u>ርር እ ር</u> ሞ ርሞ	CCTCCTCCTT	TT A ATATOAT	GTGAGGATTT	GTTACTGG
195	TGTGACCAGTCT ACACTGGTCAGA	CGGACGAGGA!	TGATATTAA	CACTCCTAAA	CAATGACO
5	11	21	31	41	
•	A A CA GTA CATGG	AGGCCTGA CCT	TGTGGGGGC	ACAGGGTGGA	ACCTTAGO
200	1 AACAGTACATGGA TTGTCATGTACC	rccggactgg <i>i</i>	ACACCCCCG	TGTCCCACCT	TGGAATCG
. 5	61	71	81	91	
	_ TGAATATAGTGTC	STGTCTCAAGA	\GGAAGTCAG	GGTACTAGCT	CAGTGCTC
205	TGAATATAGTGTC ACTTATATCACAC				GTCACGAG
5 '	11	21	31	41	
	AATCTCCAGGTAC	CTATATATACA	TTTGCCCGT'	TTTATCTC TA	ATGTGAAA
210	AATCTCCAGGTAC TTAGAGGTCCATG	TATATATGT	'AAACGGG CA	•	
5 '	61	71	81	91	
	TAAATCCCCAAAC	ACTTGTTTAT	'CGTGTAG CG'	racctaaaag:	ACTATTCT
215:	1 ATTTAGGGGTTTG	ITGAACAAATA	.GCACATCGC	ATGGATTT TC	IGATAAGA
5 '	11	21	31	41	
	ATTATGGGTGTCC	CCACTTTCTT	GGTTTGG TC!	ACCCCGAT CC	CCGGTCT
220	TAATACCCACAGG	GGTGAAAGAA	CCAAACCAGI	rggggctagg(3GGCCAGA
5 '	61	71	81	91	
_	TCTGCTGTATCTA	GAACAGTGAC	TATAAATGAT	GTATGGGAA!	CAGTGTTT
2251	AGACGACATAGAT	CTTGTCA CTG.	ATATTTACTA	ACATACCCTTA	\TCACAAA
5 '	11	21	31	41	
	CCATATGATCTGT	TGTCTGGAGT	ATATGCTACA	TGTTCAATT!	\CTGTACA
2301	CCATATGATCTGT GGTATACTAGACA	•		CACAAGTTAA	rgacatgt
5 '	61	71	81	91	
	AAAACCCAGTGCAG	GCTGATGATG	CAAAGCAGTC	CTCTCTCTGT	TACAGTG:
2351	TTTTGGGTCACGT	CGACTACTAC	GTTTCGT CAG	GAGAGAGA CAC	ATGTCAC
5 1	11	21	31	41	
	CCCCACCTATTTA	AAAATCA CGTA	ACAASCC CAG	AACACTGTGF	AACACTT
2401	CCCCACCTATTTAA GGGGTGGATAAATT	TTTAGTGCA	rGTTSGGGTC	TTGTGACACI	'TTGTGAA
5 '	61 AACATAAGAACAAA TTGTATTCTTGTTT	71	81	91	
	************	ACGCAGCGTCT	GGATTCTTT	CCAAGGAGAG	CAGCTTT
2451	TTGTATTCTTGTTT	GCGTCG CAGA	CCTAAGAAA	GGTTCCTCTC	GTCGAAA
5 1	11	21	31	41	
_	CTCCACAGGAACAC	AGTAACAAAA	GAGGTC CGC	CGCCATCCAC	ACCCAGC
2501	GAGGTGTCCTTGTG	TCATTG TTTT	'CTCCAGGCG	GCGGTAGGTG	TGGGTCG
5 '	61	71	81	91	
_	CAAGACACCTCAGA	GGCCAT AGGG	ACAACCTCC	TTGCTGG CCA	ACACCTG
2551	GTTCTGTGGA GTCT	CCGGTA TCCC	TGTTGGAGG.	AACGACCGGT	TGTGGAC

5 '			31	41
2601 CTGG	AGCAGGGGCAC	AGGTC CCAG	CAACTGATCCT	CAGTGGATGGGTCC
ZOOT GACC	TCGTCCCCGTG	TCCAGGGTCG	STTGACTAGGA	GTCACCTACCCAGG
5'	61	71	81	91
CAGT	CAAAGCCTTAA	TGGGCTCTCT	TTTGAAGGGG	A A A C A A C A A C A A A A A
2651 GTCA	GTTTCGGAATT.	ACCCGAGAGA	AAACTTCCCC'	TTTCTTTCTTAAAG
·				
5'		21	31	41
2701 AGCT	TATGATATCCA TACTATAGGT	ארבאיד ארדים מידע מידים מידים	AGTTGATGAGT	TAGTAAATTCCAA ATCATTTAAGGTT
1 CGA	1171011111001	CIMINAIN	1 CANCIACIC	MICHILIMAGGII
5 ¹	61	71		
2751 AAAAA	AAGATGATTT	TATAT GTATG	ACATAAAAAAA	ATCTTTGTAAAGT
TTTTT	TTCTACTAAAA	TATA CATAC	TGTATTTTTT	TAGAAACATTTCA
5 '	77	21	31	41
	GTGCAATAATI	TAAA GAGGTO	CTTATCTTTGC	ኋኋ ኒልጥጥጥልጥልልልጥጥልጥኒ
2801 GCGTT	CACGTTATTAA	ATTTCTCCAC	GAATAGAAACG	TATTAAATTTTA CATAATTTATAAAAT
	~ ~			
5 '	61	71		91
2851 AATAT	TGTACATGTGT ACATGTACACA	GTAATTTTTC	AIGIAIICAI TACA TAACTA	TTGCAGTCTTTGTA AACGTCAGAAACAT
TIMIN	i cili cili cili cili cili			MACG I CAGAAACA I
5'	11	21	31	41
2901 TTTAA	AAAAA CTTTAC	TGTTATGTTT	GTATAATAGA	ACATTAATCATTTA IGTAATTAGTAAAT
AAA'I"I".	L'I'I'I'I'GAAATGA	ACAA TACAAA	CATATTATCT	I'G'I'AA'I"I'AGTAAAT
5 '	61	71	81	91
20E1 TTATAA	CTCAGACAAG	STGTAAATAA	ATTCATAATT	CAAA CAGCCAGTAT
AATATI	GAGTCTGTTCC	ACATTTATT	TAAGTATTAAC	STTTGTCGGTCATA
5'	11	21	31	41
		ያገግል CAጥፕርርን	AAAAATCTCTZ	ݖݽᢕݽݽݽݵݽݽݽݾݽ ┱┸
3001 TATACG	TATATACCCAC	AATGTAACG	TTTTAGAGAT	ATCTTTGTTCTATT AGAAACAAGATAA
5 '		71	81	91
3051 CACATG	CTTAAAGAAGT Caammmeenee	AAGAAATCTT TTCTTTDGDZ	LTTG TGGATAT	GTAATTATACATA CATTAATATGTAT
GIGIAC	OAMI I I CI I CI	TIÇITINGAF	michecinin	CALIMIMIGIAL
5'	11	21		41
3101 TAAAGTA	ATATATATATG'	TATGATACAT	GAAATATATT	TAGAAATGTTCAT
ATTTCA!	PATATATATAC	ATACTATGTA	CTTTATATAA	ATCTTTACAAGTA
5 1	61	71	81	91
3151 AATTTTA	ATGGATATTCT	TTGGTGTGA	ATA ATTGAAT	ACA ACATTTTA A
LAAAATT TETE	TACCTATAAGA	AACCACACT	TAT TAACTTA	TGTTGTAAAAATT
5 '	11	21	21	4.1
3 201 AATGAAA		21 .AAAAAAAA	31 ממממממממ	41
2244 227 6244	" " " " " " " " " " " " " " " " " " "	~ ~ 3 ~ 3 ~ 3 ~ 3 ~ 3 ~ 3 ~ 3 ~ 3 ~ 3 ~		



15/41

Figure 12



5 ¹	61	71	81	91	
CEI ATT	TATATTG TTTGC	CTTTCTAAA	GTTTCTTCTAA	ATGTTCCATC	GTC
TAAT TO	ATATAACAAACGI	GAAAGATTT	CAAAGAAGATT	TACAAGGTAC	CAG'
- 1	11	. 21	21	41	•
5'		יייא יייייי כיכיכיייאי	31 		ייייבירי א
701 777 7	TAAAAAA TATACA TTTTTTATATGT	TATT GGCTA	A ATTT A AGCAG	ATTCACCCC	יבטטבי
1111	111111111111			•	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
5 '	61	71	81	91	
JE1 GAGA	TAGCTCAGAGGT ATCGAGTCTCCA	TAAGAGCAC:	CGACTGCTCTT	CCAGAGGTCC	TGAG
OTCT CTCT	'ATCGAG TCTCCA	ATTCTCGTG/	ACTGA CGAGAA	GGTCTCCAGG	ACTO
. .	11	21	31	41	
5 '		4 L (1) (1) TOCTOO	יטייטא מאמימיטאי ביייטיטא מאמימיטאי	ェエ アピヤピ ጥ カ ጥ カ ご	איייא ריי
801 TTCA	ATTCCCAGCGAC TAAGGGTCGCTG	CACAIGGIGG	CICACAGCCA:	CIGIAAIAG OTATTA DADA	$\Delta T A C$
AAGI	TANGGG I COCTO	GIGIACCACC	ONGI GI COCI.		
5 ¹	61	71	81		
GATC	TGACGC CCTCTT(ACTGCG GGAGAA(CTGGAGTGTC	TGAAGACAGC	CACAATGTAC	TCAT
851 CTAG	ACTGCGGGAGAA(GACCTCACAG	ACTT CTGTCGI	ATGT TACATG	AGTA
			2.7	4.7	
5'	11	21	31	41	
901 ATAT	ATTAAATAAATTA TAATTAAT	YTATTAGAAA ramaamoonn	ATTUTTUTAAC	TGTATCATT ממידאמיי מסמי	አጥአጥ TATA
TATA	TAALLIALLIAL	IAIAAICIII	IAAGAAGAIIC	MAIDAIAN	THI
5 '	61	71		91	
GAATZ	ATTTAA TATATA	AGTAAATGC	CTCAGGAAATA	TAAACTTGG	TTAA
951 CTTA	ATATATATAATTTA TTATATATTAAA1	TCATTTACG	GAGTCCTTTAI	'ATTTGAACC'	AATI
5'	11	21	31	41	7000
1001 AAATO	CAAAGA ACTTCAT STTTCT TGAAGTA	GAGTAGTGG	GCCA CAAAAAA GCCT CTTTTT	TGTGTACCAC	3000
TTTAC	FITICI IGAAGIA	CICALCACC	CGGIGIIIII	ACACATOOT	
5 '	61	71	81	91	
AAGAC	CGGAG GGAGGGG	AGA AGGAAG	GATGGAGATA	GAATTTTGC	CTCT
1051 TTCTG	CGGAGGGAGGGG GCCTCCCTCCCC	TCTTCCTTC	CCTACCTCTAT	CTTAAAACG	AGA
5'	11	21	31	41	700 Z
1101 GCATT	CCTTGGGCTGGC GGAACCCGACCG	ACAGGTATA!	TIGCTGTGGGA	MIIGGGAAAC	IAC
CGTAA	GGAAC CCGACCG			imiccollic	,,,,,,
5 '	61	71	81	91	
<u>አ</u> አርርአ	AGCTG CAAAGCT(GG CGGAACI	CGTTTCCGCA	AGCTGGGCTC	ATC
1151 TTCCT	TCGACGTTTCGAC	CCCGCCTTGA	GCAAAGGCGT'	rcgacccgae	;TAG
5'	11	21	31	41 rma a a a cama	idi Cim
1201 TAAGT	GTCCATGCATGGC CAGGTACGTACCC	JIG CCACACI ZNO COTOTON	CCAGTGAACT CCTCDCTTCD	₹₩₩₩₩₩₩₩ ₽	TGI
ATTCA	CHOOM LEGINCE	SACGGIGIGN	COL CACLLOM	mar ratorne	~ ~~~~
5 '	61	71	·81	91	
משייים כי	AGAGA TGTAGAGA	TGCTCACAA	TAG TACAAAG(CG GGAGGGA	.GGT
1251 07700	POTOT A CATOTOT	ימרכמכיזכייי	ልጥሮ ልጥርጥጥጥ ሮ ር	っぱっしんしょう	CCA


5 '	11	21	31	41 TTAAA CATCTGCAT ATTTGTAGACGTA	
TTA LOCA	TCCAGACTAAGA	GAAG AAAA	CCATTGCTGAT	TAAA CATCTGCAT	7
1301 TAA	AGGTCTGATTCT	CCTTCTTTTT	GGTAA CGACTI	ATTTGTAGACGTA	7.3
5'	61	71	81	91	
ጥር እ	CCCCCCCACCTC	יב ירמידמ רמרמרי	סב סב	ירט הע הערע הערע ארג. פון	٦,
1351 ACT	CGCGGGGGTGGA	GTAT GTGTG'	TGTGTGTGTGT	CACACACACACAC GTGTGTGTGTGTG	ۍ ۲٦
		•		• •	_
5'	11	21	31	41	
1401 CAAC	CCAAACAGAACAA	ATACACATGO	CATGTCTACAG	CCTGCAGGAACAA GGACGTCCTTGTT	A
GTTC	GTT1G1C11G1,1	TATGTGTAC	FIACAGATGTC	GGACGTCCTTGTT	Ί.
5'	61	71	81	91	
ATGO	TATGTCTGTGAG	GAAC CAGGAC	SATGCACAGGT	CCTAACCTCTGTC GGATTGGAGACAG	т
TACC	CATACAGACACTC	CTTGGTCCTC	TACG TGTCCA	GGAT TGGAGACAG	A
- .		21	2.7	4.7	
5'	11	TAMA AMAN AA	31	41	~
1501 CCTA	CAAGCCCTGAAG CTTCGGGACTTC	ICIGGICAGG	CARATGIA	CAAAAGCAGGCTA. GTTTTCGTCCGAT	H
5'	61	71 .	81	91 PAGGAACAACCTA PTCCTTGTTGGATA	
1551 GGAA	GCTGTTTAGTGA	AAGA TTTTTT	TCTT CAACTC	raggaacaaccta:	r
TOOT CCTT	CGACAAATCACT".	ITCTAAAAAA	AGAAGTTGAG	ATCCTTGTTGGATI	Ŧ
5 ' ·	11	21	31	41	
TTCC	TAGGAT TTGGAGA	GTG CTCAGG.	AGGAAACATTO	CAGA CAACTGATGO	3
1601 AAGG	ATCCTAAACCTCT	CACGAGTCC'	TCCTTTGTAA(CAGA CAACTGATG(STCTGTTGACTAC(3
		eg g	0.1	0.7	
5 ' mamar	11 O C C C A C A	1 1	иссе стусти В Т	91 GTTGTGCTCATGT CAACACGAGTACA	,
1651 AGAGA	I G I G I A C C C CAGA A C A C A T G G G G T C T	'AAGTCCATA	CCCCATCAAI	CAA CACGAGTACA	
1101101					
5'	11	21	31	41	
1701 ATGTO	CTAGA TATATTA	GCA CAGCCTO	SCCTTCTGCTG	CACAACGCCTTAG GTGTTGCGGAATC	ţ
	•				
51	61	71	81	91	
AGACC	CGGCCTTTCAAT	GAG CTTAGCT	TGTGCTCTGT	TTCTGCTCTCTTA AAGACGAGAGAAT	
TOTGG	GCCGGAAAGTTA	CTCGAATCGA	ACA CGAGACA	AAGACGAGAGAAT	•
	11	0.1	2.3	41	
5'	እ እ እ ሮሞ እጥሮሮጥሮጥ L T	ገን ር፤ ሙሙሙሙን ን ሙ ጉ	אמא ארא א אמי. אמא ארא א אמי	41 TATGCATCTTGCC ATACGTAGAACGG	
1801 CCAGA	TTTGA TACCACAO	TCAAAATTA	TCTTGTTTTC	ATA CGTAGAACGG	
00.1011					
5 '	61	71	81	91	
1851 TTGGC	TTGAG CCTTTTCC	TTTTCAATG	CTGACTTCTC	CCCTTTCTCTCCT	
AACCG	AACTCGGAAAAGC	AAAAGTTAC	SACTGAAGAG(eggaaagagagga	
51	11	21	31	41	
GTGCTC			AGGGACAACTT	TTAAGGAGGCGT	
				AATTCCTCCGCA	

5'	61	71	81	91
1051 GTCCCTG	GTAGGGGC	ATCCCTGTTCA	CCAGGTGCC:	91 IGTCATCACCCCACT ACAGTAGTGGGGTGA
CAGGGAC	CCATCCCCG	TAGGGACAAGT	GGTC CACGG	ACAGT AGTGGGGTGA
5 '	11	21	31	41
ТСАСТСА	CATCTACC	CTGGTGACTAT(GGGTTCCTCT	TTOTTTOTT
2001 ACTGACT	GTAGATGG	GACCA CTGATA	CCCAAGGAGA	TTGTTTGTAGGGAAC ACAAACATCCCTTG
5'	61	71	81	91
2051 GGTGGCT	CCAGGTGG	AGGCATCAATC	TGTTGGGTTC	CTGGTTCCCGGCTGC SACCAAGGGCCGACG
CCACCGA	GGT CCACC.	ICCGTAGTTAGA	CAACCCAAG	BACCAAGGGCCGACG
5 '	11	21	31	41.
CTTTGGT	TTTGAAAG:	rctcttctctg:	CATATTCCTA	CCCTGCATTTGCTT
2101 GAAACCA	AAA CTTTC!	AGAGA AGAGACI	YATAAGGAI	CCCTGCATTTGCTT CGGACGTAAACGAA
·	61	77	0.7	91
5 '	61 maamaamaa		81	ON THE REPORT OF THE
2151 7G7G7GG	TGC TGATGC	LIGIG CGCAGCA	CCTAAGAAC	GATGACTCTCCATC CTACTGAGAGGTAG
ACACACC	Aconomic			
5 '	11	21	31	41
AGTCACAC	GACTCCCCC	TGTTGCAAAGT	GTCAGGCTG	ACTCGACAGTCACC TGAGCTGTCAGTGG
TCAGTGT	CTGAGGGGG	ACAA CGTTTCA	CAG TCCGAC	TGAGCTGTCAGTGG
5 '	61	71	81	91
TAAAATO	CTGAGTCAG	TCACACACAGG	CTGTCAGCC	ACGGCTTCCACTTG
2251 GTAAAATC	BACTCAGTC	AGTGTGTCC	GACAGTCGG	ACGG CTTCCACTTG TGCCGAAGGTGAAC
			•	
5'	11	21	31	
2301 CATGGCTA	ayy Gywy y y M.T.C.T.Y.T.T.	TCACACGTGAG NGTGTGCNCTC	TTT CTGTTG	CTGGCTGACT GACCGACCGACTGA
GIACCGAI	AAGAIAAA	AGIGIGCACIC	annon canto	011000110001101011
5 '	61	71	81	91
GGCATTAT	CTATGCTA	AGTTGAAATCA	GGGGTGCCC	AGCAGAGCCCATCA CCGTCTCGGGTAGT
-5 t	11	21	31	41 CCGATGAACGTATT
	GTCTTTGA	AACAAAGCTGT	ACGGTTTGA:	CGATGAACGTATT
2401 AAGAGTGA	CAGAAACT	TGTTTCGACA:	rgc caaact?	AGCTACTTGCATAA
5'	61	71	8T	91 GAAGGCAGGCTGT CCTTCCGTCCGACA
2451 TAAAGCAT	TT CATGCAA N N GTN CGTT	TGA CAAAGTGC TA CTCTTTCA CC	TCAGIAGIO	CTTCCGTCCGACA
ALLICGIA	MAGINCGII	ACIGITICACC	Morania	
51	11	21	31	41
2501 GACCAGTC	TGCCTGCTC	CTTACTATAA1	TGTGAGGAI	TTGTTACTGGAAC
CTGGTCAG	ACGGACGAG	GAATGATATTA	ACACTCCTA	AACAATGACCTTG
5'	61	71	81	91
አ ረመአ ረአ ምርረ				GAA CCTTAGCTGA
2551 TCATGTAC	CTCCGGACT	GGAACACCCC	GTGTCCCAC	CTTGGAATCGACT


	5 ¹	11	21	31	41
26	O1 ATATAGTGT	GTGTCTCAA	GAGGAAGTC	GGGTACTAGC	rcagtgeteaat Agteaegagtta
	TATATCACA	CACAGAGTT	CTCCTTCAGI	CC CATGATCG	AG TCACGAGTTA
!	5 '	61	71	81	91
0.0	CTCCAGGTA	CTATATATA	^ጉ ል ጥጥጥር ር ር ር ር	ያምምምምልምምም የ	א אייי א א מבייבידי א
26	51 GAGGTCCAT	GATATATAT	TAAACGGGC	AAAATAGAGAT	TACACTTTATT
		•			
į	5'	11	21	31	41
27	01 TACCCCAAA	CACTTGTTT	AT CGTGTAGC	GTACCTAAAAG	A CTATTCTATT TGATAAGATAA
	IAGGGGIII	3 I GAACAAA)	AGCACATCG	CAIGGATTTC	TGATAAGATAA
9	5 t	61	71	81	91
27	ATGGGTGTC	CCACTTTCI	TGGTTTGGT	CACCCGATCC	CCCGGTCTTCT GGGCCAGAAGA
21.	TACCCACAGO	GGTGAAAGA	ACCAAACCA	GTGGGGCTAGG	GGGCCAGAAGA
_			0.5		
5		11	21	31	41
280	O1 GCIGIAICIA	IGAACAGTGA TOTTOTOTOTO	CTATAAATG/	ATGTATGGGAA	41 TAGTGTTTCCA ATCACAAAGGT
5	t	61	71	81	91 ACTGTACAAAA TGACATGTTTT
201	TATGATCTGT	'TGTCTGGAG	TATATGCTAC	CATGTTCATTT.	ACTGTACAAAA
201	² ATACTÁGACA	ACAGACCTC	ATATACGATO	TACAAGTAAA'	TGACATGTTTT
5	•			24	
5		にしむなかむなかれた エエ		31	
290	1 TGGGTCACGT	CGACTACTA	CGTTTCGTCA	GAGAGAGACA	41 GTACAGTGCCC CATGTCACGGG
5		61	71	81	91
295	1 CACCTATTTA GTGGATAAAT	AAAATCACGT	ACTTGCCCA	GAACACTGTG	AACACTTAAC
	GIGGATAAAT	TTTTAGTGCA	ATGAACGGGT	'CTTGTGACACT	TTTGTGAATTG
5	t	11	21	31	41
200			TGGATTCTT	TCCAAGGAGAG AGGTTCCTCTC	CAGCTTTCTC
300	$^{\perp}$ TATTCTTGTT:	rgcgtcgcag	ACCTAAGAA	AGGTTCCTCTC	GTCGAAAGAG
 .	é				~
5 ') ************************************)	71	81 3333377337	91
305:	CACAGGAACA (AGTAACAAA TYCDYYCYYYY	TCTCCDGGCC	81 CCGCCATCCAC GCCGTAGGTG	PCCCAGCCAA
	01010011010	J1 @11 1 G1 1 1	rereasou	SCCCCIACCIO	1000100011
5 '	1		21	31	41
3101	GACACCTCAGA CTGTGGAGTCI	GGCCATAGG	GACAACCTC	CTTGCTGGCCA	ACACCTGCTG
5203	CTGTGGAGTCT	CCGGTATCC	CTGTTGGAG	SAACGACCGGT	TGTGGACGAC
5 1	6	1	71	81	91
_	_				
3151	GAGCAGGGGCA CTCGTCCCCGT	GTCCAGGGT	CGTTGACTAG	GAGTCACCTA	CCCAGACGTC
5 '	1:		21		41
3201	CCAAAGCCTTA. GGTTTCGGAAT	ATGGGCTCT (CTTTTGAAGG	GGAAAGAAAG	AATTTCAAGC
	COLLECTORAL.	LLLCCCGAGAC	WARACA T T C C	COLLECTION.	TAMMATICA

		61	71	81	91
5 '	TTATGATATC			AGTTAGTAAA	TTCCAAAAAA
3251	TTATGATATC AATACTATAG	ת מיית מיית מיית או	TATCAACTAC	TCAATCATTT	AAGGTTTTTT
J2J2	AATACTATAG	GITATAATAA	IMICALICALIC		
		11	21	31	41
5 '	AAAAGATGAT		mananna a a a	እ እ አ ልጥርጥጥጥ ር	TAAAGTGCGC
2201	AAAAGATGAT TTTTCTACTA	TTTATATGTA	LACATAGA	TTTTAGAAAC	ATTTCACGCG
3301	TTTTCTACTA	AAATATACAT	ACIGIATITA		
		<i>~</i> 3	71	81	91
5 '	AAGTGCAATA	61	▗▗ ▗▗ ▗▗ ▗▗	・TGCATTTATA	TAAATATTAA
2251	AAGTGCAATA TTCACGTTAT	ATTTAAAGAG	GICIIAICII	ACGTAAATAT	ATTTAATATTTA
3331	TTCACGTTA'I	TAAATTTCIC	CAGAATAGA	110011	
			21	31	41
5 '	ATTGTACATG	11	~~ mama ma	CARTTCCACT	CTTTGTATTT
2401	ATTGTACATG TAACATGTAC	TGTGTAATTT	TTCATGIAII	GTAAACGTCA	GAAACATAAA
3401	TAACATGTAC	ACACATTAAA	AAGTACATAA	GIZHHIOOL	
			71	81	91
5 '		61		ነ አ ረን አ ረን ምጥ <u>ል</u> ል	מדים בידים בידים
	AAAAAAACTT TTTTTTTGAA	TACTGTTATG	TTTGTATAAT	TOTAL ATT	AGTAAATAAT
3451	TTTTTTGAA	ATGACAATAC	AAACATATTA	ICIIGIA	
			~ =	31	41
51		11	21	Σ	ссастатата
	TAACTCAGAC	AAGGTGTAAA	TAAATTCATA	$\mathbf{A}\mathbf{I}\mathbf{I}\mathbf{C}\mathbf{A}\mathbf{A}\mathbf{A}\mathbf{C}\mathbf{C}\mathbf{C}$	GGTCATATAT
3501	TAACTCAGAC ATTGAGTCTG	TTCCACATTT	ATTTAAGTAT	IMAGILIOLO	
	•			0.1	91
5'		61	/ <u>T</u>	ᠳᢕᠬ᠉ᡎ᠘ᡴᠳᡎᢙ	TATTATTCAC
	TGCATATATG	GGTGTTACAT	TGCAAAAATC	TOTATOTATO	AAGATAAGTG
3551	TGCATATATG ACGTATATAC	CCACAATGTA	ACG1"1"1"1"AG	AGAIAGHMIC	
			. .	27	41
5 '				አመአጥረሞአ አጥባ	АТАСАТАТАА
	ATGCTTAAAG.	AAGTAAGAAA	TCTTTTGTGG	AIAIGIAAI	TATGTATATT
3601	ATGCTTAAAG. TACGAATTTC	TTCATTCTTT	AGAAAACACC	IAIACAIII	
				0.1	91
5 '		· -		$m_{\lambda} m_{UU} m_{\lambda} C \lambda \Delta \Delta$	TGTTCATAAT
2651	AGTATATATA' TCATATATATA	TATGTATGAT	ACATGAAATA	ATAAATCTTT	ACAAGTATTA
3651	TCATATATAT	ATACATACTA	TGTACTITAL	WIIMMIT O'T	
			21	31	41
5 1	TTTAATGGAT		~~~~~~~~~~	CAATACAACA	TAAAATTTTT
2701	TTTAATGGATA AAATTACCTA	ATTCTTTGGT	GIGNATAWII	CTTATGTTGT	ATTTTAAAAA
3701	AAATTACCTA'	TAAGAAACCA	CHCITHITH	01111	
			71	81	91
5 ') KAAAAAAAAA TTTTTTTTT	61.	/ T ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	ΔΔΑΩΤΤΤΤΤ	TTTTTTTTT
2751	AAAAAAAAA	AAAAAAAAA.	ᡎᠬᠳᡎᡎᡎᡎᡎᡎᡎᡎ	AAAAATTTT	AAAAAAAAA
3/51	TTTTTTTTTT	L.J.,J.,J.,J.,T.,T.,T.,T.,T.	111111111	1111111	
		:	o 1	21	41
5 '	_		▗▗ ▄▗▄▄▗▗▗▄╓▞ᡢᡏ	ጥአ አ ୯୯୩୩GAA	GGGCAGGCAA
3801	TTATTCCAGA(AATAAGGTCT(JATTAAAGACA	TO A TO CALL OF THE CALL OF TH	ATTGGAACTT	CCCGTCCGTT
200T	AATAAGGTCT(TAATTTCTG	IGHICINGAA		
	_		71	81	91
5 ') ACGCTCGCAL CTCCACCTTC	51	ᄼᆂ ᇽᄽᄱᄽᄶᄽᄽᄺᅲᄼ	AGGGACCATT	TTCTTCTTGA
2051	GAGGTCGGCA	ATGUTGTUAA	ℳℷ℞℧℞℞℧ℷℂ	TCCCTGGTAA	AAGAAGAACT
202T	<u> </u>	ACGACAGTT	2 THICIICHG		

5 '		11	21	31	41 GCTCCGGAAT CCGAGGCCTTA
3901	ACATGCAGTC	ACTTTCCTGA	TTGCTCTTC	A CATCCTCAAG	GCTCCGGAAT
3901	TGTACGTCAG	TGAAAGGACI	'AACGAGAAG'	TGTAGGAGTTC	CGAGGCCIIA
. 5 (61	71	81	91
_	TCCGGGGGTG	TGGTGGGCTT	TGATCTCAG	GACTCTGGAGG	CAGAAGCAGG
3951	AGGCCCCCAC	ACCACCCGAA	ACTAGAGTC	CTGAGACCTCC	CGTCTTCGTCC
5'1		11	21	31	41
_	CAGATCTCTG	TGAATATGAG	GCCAGCCTG	CACTACACAGA	GCTCCAGACC
4001	GTCTAGAGAC	ACTTATACTO	CGGTCGGAC	GTGATGTGTCI	CGAGGTCTGG
_ ·		61	71	81	91
5.	maaaa	አ ረን ሞረን ሞረን አ	አሮርርጥርጥርጥር	CAAAAAGAAAA	TAAAAACTGT
4051	TCAGTACCGA	TGTAGTACTT	TGGGACAGAG	TTTTTCTTT	ATTTTTGACA
		- -	21	31	41
5',	mamammam Am A	11 CCDTDGTGTT	ል ል ል ርጥር ል G ል G	TCTGAGTAAT	'GTCGGGCTGA
4101	ACACAAAGAT	GGTATCACAA	TTTGAGTCTC	CAGACTCATTA	CAGCCCGACT
- ,	,	61	71	81	91
5			N COMMON COT	TTGACGAGGC	GCTGAACAGT
4151	GTACGAGCCC	ACAAATTGTA	TGGAAGTCGA	AACTGCTCCG	CGACTTGTCA
5 '	-	11	21	31	41
_		CTTCCCCAC	CGGTGGCTGT	GTTTGTGCTC	AAGTCCACCG
4201	GTTTCAGACC(GAACCCCTC	GCCACCGACA	CAAACACGAG	TTCAGGTGGC
5 1	•	51	71	81	91
4051	TGAAATCCTG <i>I</i> TGAAATCCTGAC	TTGTGAATT'	TGGACAACCG	TGTCCTTCTT	CTTGGCCTTC
425I	ACTTTAGGACI	'AACACTTAA.	ACCTGTTGGC	ACAGGAAGAA	DAACOOJAAO
5 '	3	L 1 :	21	31	41
-	CATGCAACCT	CAACTTCAT	GTTGGTCATT	TTGTCAAAAC	ACTGTGTGAT
1301	CATGCAACCTC GTACGTTGGA	GTTGAAGTA	CAACCAGTAA	AACAGTTTTG	TGACACACIA
Б.I	6		71	81	91
:_ (ammonora mara	מיז מיז כיניכיכי	ለምምርርልርልምል	TGTAGAGATG	TAGTCTGCCT
1351	GTTTTTATCAA CAAAAATAGTT	'ATATGACGG'	PAAGGTGTAT	ACATCTCTAC	ATCAGACGGA
51	1	.1 2	21	31	41
	1 GGCTTTCCTTT CCGAAAGGAA	TCTTTAGCCA	AATCGAATGC	TCTTGATCAT	GCCCTCAATC
FAOT (CCGAAAGGAAA	AGAAATCGGT	TAGCTTACG.	AGAACTAGTA	CGGGAGTTAG
5 '	6	1 7	71 .	81	91
_	o CATCTCTAGC	TTTTATCACO	TCTCTGCTA	ATTCCTGAAA	CTTGAATCGA
451	AGTAGAGATCG	AAAATAGTGC	'AGAGACGAT'	TAAGGACTTT.	GAACI TAGCT
5 '	1	1 2	1	31	41
	᠘ᢗᡆᠬᠳᠬ᠇ᡎᡳᡎᡎ	GGTTCATCTC	'AATGGTGAT	GTTCAGTTCC'	TTCTGAATCT
5017		ሮሮሽ አርሞአርአር	ጥጥአ ሮሮኔ ሮሞል (CAAGTCAAGG	AAGACTTAGA



				4 4
5 '	11	21	31	41
E201	CTATAAȚCAA TGTAC	SACTGA TAATA	TTATG CCAGC	AAATAGTTTTGAAGT
5201,	GATATTAGTTACATO	TGACTATTAT	AATACGGTCG'	TTTATCAAAACTTCA
5'	61	71	81	91
	CCTAGGCACAGTGGG	AGGAGGTTTT	GTTCCACGCT	GTTCATAAGCCAATA CAAGTATTCGGTTAT
5251	CATCCGTGTCACCC	TCCTCCAAAA	CAAGGTGCGA	CAAGTATTCGGTTAT
`	JOHI CCCIOI CITTO			
5 '	11	21	31	41
ر د	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	יייים או או מכלא כיאו	א כתיויכות א איתיי	rggga Cattcacatc ACCCTGTAAGTGTAG
5301		7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	$\mathbf{r}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$ $\mathbf{n}_{\mathbf{C}}$	ACCCT GTA AGTGTA G
,	366616611116166	AATTICCIGI	IGAACHIIII	.ccci ciii.ciciqiAG
- .	61	71	81	91
5'	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~			
5351	rgreererrearers	ATCIGGCTCCC	AGIGICACIO	CTCTAACACGGTCCT SAGATTGTGCCAGGA
JJJI	ACAGGAGAAG TAGAC	TAGAC CGAGGC	STCACAGIGAG	SAGAT TGTGCCAGGA
		21	2.1	4.1
5 '	11	21	31	41
E401 T	'AGAGGGACAATTTA'	rcccrgccrc:	GCTTGATCTI	TATGCATGTATCTGT TACGTACATAGACA
2401.7	TCTCCCTGTTAAAT	AGGGA CGGAGI	ACGAA CTAGAA	TACGTACATAGACA
	•			
5' .	61	71	81	91 .
A	TTCTTCCAGCCATC	CCTGG CGACCI	GATTTTTCTA	AGGCACCCAAAACT TCCGTGGGTTTTGA
5451 _T	'AAGAAGGTCGGTAG(GACCGCTGGA	CTAAAAAGAT	TCCGTGGGTTTTGA
5'	11	21	31	41
. G	TAAGCTACT TCTTAT	CAATCTATAAT	TCTGAGCATA	TTAGTTAGCCTGAG
5501 🖰	TAAGCTACT TCTTAT ATTCGATGA AGAATI	TTAGATATTA	AGACTCGTAT	AATCAATCGGACTC
5 '	. 61		81	91
	CTCCAGGATATCTTT GAGGTCCTATAGAAA	CTTC CCTATA	CTCAGTCCAG	TTTTAGCTGCCCAG
چ 5551	CACCTCCTATAGAAA	GAAGGGATAT	GAGTCAGGTC	AAAATCGACGGGTC
J	01.001.001.1			
5 '	11	21	31	41
70	A CCA TTCA A ACCTGA	TOTA CGAGTA	GATCACTCCT	GTCTACAGCTTGTT
5601 🛱	TCCTAAGTT TCGACT	AGATGCTCAT	CTAGTGAGGA	CAGATGTCGAACAA
_	1CCIAAGI į I COIICI	110111 001 011	011101011	
5 '	61	71	81	91
5.		7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	,	
5651 C	CAGATCTTGTTTCTC STCTAGAACAAAGAG	₩₩ĠĊ <i>Ċ</i> ĊŢĠĠ₩	rCCCTACTCC	CHOO THICHITCHA
G	3TCTAGAACAAAGAG	IICGGGACCI.	ICGG IAGICO	orcentre in the contraction
	11	21	31	41
5'	AACAATCCCTTTCTA	2 T		
5701 AA	AACAATCCCTTTCTA. FTGTTAGGGAAAGAT	ATCATGGGTG.		COUNTY COCCOORAL
TJ	I"I'GI"I'AGGGAAAGA'I".	PAGTACCCACA	TCCGGGTTTC	ACTIACCOGCCITA
	C3	77	81	91
5'	61	71	ΩŢ	3.
5751 TG				
,,,, AG	}			



25/41




Figure 14

Figure 15

1	CGCCCGGCA GGTCTGTTG				
	GCGGGCCCGT CCAGACAAC				
61	AGACCCCACT GATGGTGTG	C TGCCTTTCAG	TCCAGGAAGA	AAGAAAGGAA	GGATTCTGAG
	TCTGGGGTGA CTACCACAC				
121	GATTTGGGCA AAGCCACAT	T CCTGGAGAAG	TCTGTATACT	GATGCCAAAC	CCAAGAGCTG
	CTAAACCCGT TTCGGTGTA				
181	AGCTGCTGAT GAGGCCCAG	G GAGTAGCCCA	· CGCGCCCTGA	GCTGTTGGCT	AGCAAGGCCT
	TCGACGACTA CTCCGGGTC				
241	TCCTGCTCCA TGTGGCATG	G AAAAATTATA	TGGTTTGACG	GATGAAAAGG	TGAAGGCCTA
	AGGACGAGGT ACACCGTAC				
301	TCTTTCTCTC CATCCCAG	G TATTAGATGA	ATTTGTTTCT	Gaaagtgtta	GTGCAGAGAC
	AGAAAGAGAG GTAGGGGTC				
361	TGTGGAAAAG TGGCTGAAG	а ссалалссал	CAAAGCAAAA	GATGAACCAT	CTCCCAAGGA
	ACACCTTTTC ACCGACTTC				
421	AGTCAGCAGG TACCAGGAT	A CGAATATGCA	GGGAGTCGTG	TACGAGCTGA	ACAGCTACAT
	TCAGTCGTCC ATGGTCCTA				
481	AGAGCAGCGC CTGGACACGC	G GCGGGGACAA	CCACCTGCTC	CTCTATGAGC	TCAGCAGCAT
	TCTCGTCGCG GACCTGTGC	CGCCCCTGTT	GGTGGACGAG	GAGATACTCG	AGTCGTCGTA
541	CATCAGGATA GCCACAAAA	CCGACGGATT	TGCACTGTAC	TTCCTTGGAG	AGTGCAATAA
	GTAGTCCTAT CGGTGTTTTC	GGCTGCCTAA	ACGTGACATG	AAGGAACCTC	TCACGTTATT
601	TAGCCTGTGT GTGTTCATAC	CACCCGGGAT	GAAGGAAGGC	CAACCCCGGC	TCATCCCTGC
	ATCGGACACA CACAAGTATO	GTGGGCCCTA	CTTCCTTCCG	GTTGGGGCCG	AGTAGGGACG
661	AGGGCCCATC ACCCAGGGTA	CCACCATCTC	TGCCTACGTG	GCCAAGTCTA	GGAAGACGTT
	TCCCGGGTAG TGGGTCCCAT	GGTGGTAGAG	ACGGATGCAC	CGGTTCAGAT	CCTTCTGCAA
٠.	EcoRV	:	*********	ww	mach h man ca
721	GTTGGTAGAG GATATCCTTG	GGGATGAGCG	ATTTCCTCGA	CCATCACCCC	ACCTTAGTCC
	CAACCATCTC CTATAGGAAC	CCCTACTCGC	TAAAGGAGCI	CCCATTACCAC	ACTTCATTGG
781	AACCCGCATC CAGTCTGTTC TTGGGCGTAG GTCAGACAAG	TTTGCTTGCC	CATTGICACI	CGGTAACCTC	TGAACTAACC
	CATCCTTGAA CTGTACAGGC	AAACGAACGG	*C*CCCCTTC	TOCCTOR COC	ATCAGGAGGT
841	CATCCTTGAA CTGTACAGGC GTAGGAACTT GACATGTCGG	* ACTGGGGCAA	TCTCCGGAAG	ACGGAGTCGG	TAGTCCTCCA
• • • •	TGCAACAGCC AATCTTGCTT	CCCCETT	ACCANTACAC	CAGGTGCAGG	TGTGTAGAGG
901	ACGTTGTCGG TTAGAACGAA	CCCCAACCCA	TCGTTATGTG	GTCCACGTCC	ACACATCTCC.
	TCTCGCCAAA CAGACCGAAC	TCAATCACTT	CCTACTCGAC	GTATCAAAGA	CATACTTTGA
961	AGAGCGGTTT GTCTGGCTTG	ACTTACTGAA	GGATGAGCTG	CATAGTTTCT	GTATGAAACT
	TAACATAGTT GCCATAGACT	CTCTACTTGA	ACACATCATG	ATATATGCAA	AAAATCTAGT
1021	ATTGTATCAA CGGTATCTGA	GAGATGAACT	TGTGTAGTAC	TATATACGTT	TTTTAGATCA
	GAACGCCGAC CGCTGCGCGC	TCTTCCAGGT	GGACCACAAG	AACAAGGAGC	TGTACTCGGA
1081	CTTGCGGCTG GCGACGCGCG	AGAAGGTCCA	CCTGGTGTTC	TTGTTCCTCG	ACATGAGCCT
1141	COMOTTONO ATTROCCOAGE	AGAAGGAGGG	GAAGCCCATC	TTCAAGAAGA	CCAAGGAGAT.
1141	GGACAAACTG TAACCCCTCC	TCTTCCTCCC	CTTCGGGTAG	AAGTTCTTCT	GGTTCCTCTA
1201	CACATTTTCC ATTGAGAAAG	GGATTGCTGG	TCAAGTGGCA	AGAACAGGCG	AAGICIIGA.
	GTCTAAAAGG TAACTCTTTC	CCTAACGACC	AGTTCACCGT	TCTTGTCCGC	TTCAGAACTI
1261	CATTCCCCAT CCCTACCCCG	ACCCTCGCTT	TAACAGGGAG	GTGGACCTGT	ACACAGGCTA
	GTAAGGGCTA CGGATGCGCC	TGGGAGCGAA	ATTGTCCCTC	CACCTGGACA	16161666
	CACCACCACC AACATTCTCT	GTATGCCCAT	AGTGAGCCGA	GGCAGCGTGA	TTGGCGTGGT
_	GTGGTGCTCC TTGTAAGACA	CATACGGGTA	TCACTCGGCT	CCGTCGCACT	AACCGCACCA
					•

Figure 15 (con't)

1381	GCAGATGGTG AACAAGATCA GCGGTAGCGC	CTTCTCCAAG	ACAGACGAGA	ACAACTTCAA
2502	CGTCTACCAC TTGTTCTAGT CGCCATCGCG	GAAGAGGTTC	TGTCTGCTCT	TGTTGAAGTT
				Banit
	GATGTTTGCT GTCTTCTGCG CACTGGCCTT	GCACTGTGCT	AACATGTACC	ACAGGATCCG
1441	CTACAAACGA CAGAAGACGC GTGACCGGAA	CGTGACACGA	TTGTACATGG	TGTCCTAGGC
	CTACAAACGA CAGAAGACGC GTGACGCG	LACACHA.		
	CCACTCAGAA TGCATCTACA GGGTTACCAT	WWWWWW CONT	TCCTACCACA	GCATCTGCAC
1501	CCACTCAGAA TGCATCTACA GGGTTACCAT GGTGAGTCTT ACGTAGATGT CCCAATGGTA	COTOTTOGAA	AGGATGGTGT	CGTAGACGTG
	GGTGAGTCTT ACGTAGATGT CCCAATGGTA	COTOTTOCO.	CCACCCATCT	CCCCCACAT
1561	CTCCGAGGAG TGGCAAGGCC TCATGCGCTT	CAACCIACCA	CCTCCCTAGA	CCCCCTCTA
	GAGGCTCCTC ACCGTTCCGG AGTACGCGAA	GTTGGATGGT	TO COOTICOCA	TOTOTOTA
1621	CGAGCTATTC CACTITGACA TIGGTCCTTT	CGAGAACATG	NOCCIOGON	TOTTIGICIA
	CONCERNATE COCABACTET AACCAGGAAA	GCICITGIAC	NOOGNOOT	Monnacioni
1681	TOTAL STREET, CONCRETCE	TTTGAACTT	GAAAAATTGT	GCCGTTTTAT
	THE PART OF THE PA	AAAACIIGAA	CITITION	COCCURRENT.
1741	THE PROCESS OF A PROCESS OF THE PROC	TYCTTACCAC	AACIGGAAGC	WIRCHGI CHC
717	CATGTCTGTG AAGAAGAACT ATCGGCGGCCA GTACAGACAC TTCTTCTTGA TAGCCGCCCA	AGGAATGGTG	TTGACCTTCG	TACGTCAGTG
	· · · · · · · · · · · · · · · · · · ·			***********
	GGTGGCACAC TGCATGTATG CCATACTTCA	AAACAACAAT	GGCCTCTTCA	CAGAÇCTCGA
1801	GGTGGCACAC TGCATGTATG CCATACITCA CCACCGTGTG ACGTACATAC GGTATGAAGT	TTTGTTGTTA	CCGGAGAAGT	GTCTGGAGCT
'				
	**	CCATGACCTG	GACCACAGGG	GCTTCAGTAA
1861	The second secon	GGIVCIGO	0.00.000	
1921	CAGCTACCTG CAGAAGTTCG ACCACCCCT GTCGATGGAC GTCTTCAAGC TGGTGGGGGA	CCCCCCGCGAC	ATGAGGTGGA	GGTGGTACCT
1981	GCAACACCAC TTCTCCCAGA CGGTGTCCAT CGTTGTGGTG AAGAGGGTCT GCCACAGGTA	GGAAGTCGAC	CTTCCCGTGT	TATAGAAGAG
2041	CACCTGAGC TOCAGGAGT ACGAGCAGGT GTGGGACTOG AGGTOGCTCA TGCTCGTCCA	CCRCCTCTAG	TAGGCGTTTC	GGTAGTAGCG
2101				
2161				
2202				
2221	THE TOTAL STORY TOTAL CONTROL CONTROL	GCCAGITACA		GCTTACTATA
2221	GACACTAGAA ACGAGACACT GGTTTGATAC	CGGTCAATGT	TTTAACTGTC	
• •	East	•		*CCCCPTTCC
2221		GATGAAGAAG	CTGGGCATAC	TCCCCT1100
2281	ATATGCAGAA TTCTGGGCTG AGGGTGATGA TATACGTCTT AAGACCCGAC TCCCACTACT	CTACTTCTTC	GACCCGTATG	10000111100
	TOTAL TOTAL CONTROL GAGATGAAGT	CCCICAAGGG		
2341	ATACTACCTG TCTCTGTTCG CTCTACTTCA			
	TGTGGCCATT CCCTGCTATA CCACCTTGAC	GCAGATCCTC	CCACCCACAG	AGCCTCTGCT
2401	TGTGGCCATT CCCTGCTATA CCACCTTGAC ACACCGGTAA GGGACGATAT GGTGGAACTG	CGTCTAGGAG	GGTGGGTGTC	TCGGAGACGA
	ACACCGGTAA GGGACGATAT GGTGGAACIG GAAGGCCTGC AGGGATAACC TCAATCAGTG	GGAGAAGGTA	ATTCGCGGGG	AAGAGACAGC
2461	GAAGGCCTGC AGGGATAACC TCAATCAGTG CTTCCGGACG TCCCTATTGG AGTTAGTCAC	CCTCTTCCAT	TAAGCGCCCC	TTCTCTGTCG
	ATTCCGGACG TCCCTATTGG AGTTAGTCAC AATGTGGATT TCAGGCCCAG GCCCGGCGCC	TAGCAAGAGC	ACACCTGAGA	AGCTGAACGT
2521	AATGTGGATT TCAGGCCCAG GCCCGCGCCC TTACACCTAA AGTCCGGGTC CGGGCCGCGG	ATCGTTCTCG	TGTGGACTCT	TCGACTTGCA
	TTACACCTAA AGTCCGGGTC CGGGCCGCGG GAAGGTTGAA GACTGATCCT GAAGTGACGT	CCTGATGTCT	GCCCAGCAAC	CGACTCAACC
2581	GAAGGTTGAA GACTGATCCT GAAGTGACGT CTTCCAACTT CTGACTAGGA CTTCACTGCA	GGACTACAGA	CGGGTCGTTG	GCTGAGTTGG
	TECTTCTGTG ACTTCGTTCT TTTTGTTTTC	AAGGGGTGAA	AACCCCCTGT	CAGAAGGTAC
2641	TECTTCTGTG ACTTCGTTCT TTTTGTTTTC ACGAAGACAC TGAAGCAAGA AAAACAAAAG	TTCCCCACTT	TTGGGGGACA	GTCTTCCATG
	MODINGROAD TORROGERA			

Figure 15 (con't)

2701	CGTCGCATAT CCATGTGAAG CAGACG	GACTC	CCTGCTTGCC	GCACACACCT	CGGACAGTGA
	GCAGCGTATA GGTACACTTC GTCTGC	CTGAG	GGACGAACGG	CGTGTGTGGA	GCCTGTCACT
2761	GCAACCCAGG CTCTGCCGTG TTCAGA	ACGTC	GGCTACTCCG	TGGCTCCACC	ACTOCACCO
	CGTTGGGTCC GAGACGGCAC AAGTCT	rgcag	CCGATGAGGC	ACCGAGGIGG	ACTOGAGGCT
2821	ATGCTATTTG CTCCCAGGCC AGCACT	rgcac	TGTCTGGAGG	GGGCAGAGAC	CMCMCCMCMC
	TACGATAAAC GAGGGTCCGG TCGTGA	CGTG	ACAGACCTCC	CCCGTCTCTG	GIGICCICIC
2881	GTTCTTGCCT GCATCCTCCC ATGAGG	GTGT	GGCCAGTTCC	CTAGTTCTGT	CCCTACCACC
	CAAGAACGA CGTAGGAGGG TACTCO	CACA	CCGGTCAAGG	GATCANGACA	COGIACGACG
2941	TGCTTGGTGG CATTGGTTAG GAATGC	GACA	CACGCCCCTT	CA ACACTUCA	AATCATGTGA
	ACGAACCACC GTAACCAATC CTTACC	CTGT	GTGCGGGAA	CAACACTICA	AN CO COCCO
3001	CCTTCTTATA GGTTAACTGA GTTTGT	rGGCC	TGGACACATG	TAATGAAGGT	CACAGTCCAC
	GGAAGAATAT CCAATTGACT CAAACA	CCGG A	ACCTGTGTAC	ATTACTICA	TOTOLOGIG
3061	AGGTGACAGA GAAATCCAAA CTGTTG	ATTA	CAGGTGCACT	MCAGGIAIGC	ACARACTOR
	TCCACTGTCT CTTTAGGTTT GACAAC	TAAT	GTCCACGTGA	PACCABACCE	CECCCCECA E
3121	TATCTGGGG CACATAGGTG AGTCTG	CTCC	ACTCAGAANN	TOORINGE	CIGCCCICAI
	ATAGACCCC GTGTATCCAC TCAGAC	GAGG	TGAGTCTINN	A COMOMONOM	TON N N CONTC
3181	CCAGGGGACA CAGGGTACAT CCCAGG	CATC	GGGGAACTGA	TOCACICACI	AGTTTGGTAC
	GGTCCCTGT GTCCCATGTA GGGTCC	GTAG	CCCCTTGACT	TOTAL TOTAL OF	CCCCAATCC
3241	TCAAAGAATT AAAACACCTC CCCTCC	CCCT	CACTGTAGCC	A DECTETTEA	CCCCGTTAGG
	AGTTTCTTAA TTTTGTGGAG GGGAGG	GGGA	GIGACATOGG	TOCACCAACC	AAATCTTGTG
3301	CTTTATACAA AGAAAATAAA AGTAAG	GCAT	ATAAATTICC magggaaa	AGGTCGTTCG	TTTAGAACAC
	GAAATATGTT TCTTTTATTT TCATTC	CGTA	CHANTIANT	NTCHCHCHAT	GTTATGGCAG
3361	GGTAAAAAA AAGCATGTGA ATNNTA	LACAA (CNTCTANANI	NAGNGNCNTA	CAATACCGTC
	CCATTTTTT TTCGTACACT TANNAT	TGTT	GRAGAINI	ATACCTCATC	CTATGCCTGA
3421	AATTTAGTC ACGTCCAAAA CAAAAA TTAAAATCAG TGCAGGTTTT GTTTTT	GATT A	TA ACCTUTU	TATGGAGTAG	GATACGGACT
	AAGGCTCCAC AGCATGGCGT CCGTCT	CIAA	CCCTTCTCAT	CCGTCTCCTC	ACGGTGCAAT
3481	AAGGCTCCAC AGCATGGCGT CCGTCT TTCCGAGGTG TCGTACCGCA GGCAGA	CCCA (CCCAAGACTA	GGCAGAGGAG	TGCCACGTTA
	CAGGCAGGAC AGAGAGGAGG GCTGCA	CCCC	TACCACATTG	ACCCAGAAGG	TATCTCCTCT
3541	CAGGCAGGAC AGAGAGGAGG GCIGCA GTCCGTCCTG TCTCTCCTCC CGACGT	CCCC A	ATGGTGTAAC	TGGGTCTTCC	ATAGAGGAGA
	CORRECT CORREC	ממסטי	ATCCTGTATT	GAATAGTTCT	CTGTGTGACT
3601	CACCATTCAG ACATCCATAA GGAATG GTGGTAAGTC TGTAGGTATT CCTTAC	GGTT 1	TACGACATAA	CTTATCAAGA	GACACACTGA
	er. of				
		CCTT 7	TCCNGGGGAA	CTCTAAGGAG	TCACAGGTTC
3661		GGAA /	NGGWCCCC1 +	01101111001	
	ACACCGTGGG GATTTTCAGG ATAGCA	TCCD (CACAGAGATU	COGICGIIG	1010
3721	TGTGGCACCC CTAAAAGTCC TATCGT	ACCT (CIGICICING	0001100121011	
122.	GAGCCTTGAG AAGGAGAGAC TGACCA	GAAA C	CACTCACTCA	GCACTCTGCA	GGAGCAGGAG
3781	CTCGGAACTC TTCCTCTCT ACTGGT		GTGAGTGAGT	CGTGAGACGT	CCTCGTCCTC
	AAGATACTTT AAGATGAATC TTGGAT	AGAT 1	TTTGATACAC	CCAATACCAT	ACACACAGGA
3841	TTCTATGAAA TTCTACTTAG AACCTA	TCTA ?		GGTTATGGTA	
2001	TOTAL	ጥጥጥር (TTCCGCGCT	CTGACCCACG	GTTGTAGCGG
3901	TARGETT ACCULATE TARGET	AAAG C	GAAGGCGCGA	GACTGGGTGC	CAACATCGCC
3061	AGTGGGCTGA ACACTGTAAC ACTGTA	ሮክጥር (CATTTCCCC	AIGGGCIICI	MARKET
3961	TCACCCGACT TGTGACATTG TGACAT	GTAC G	CIMANGGGG	INCOOLLIGIT	
4021		~~ n m ~ ~	TACTGGTTA	CAAGGTGATG	TCAACAAGAG
7021	CATCTCCTCC CCTGCTGTGT CCTACTCGTAGAGGAGG GGACGACACA GGATGAC	GGTA A	ATGACCAAT	GTTCCACTAC	AGTIGITOTO

Figure 15 (con't)

4081	AAGCTATCAC AACACCAGGG C	TGTGCACAC	GTGCACACAC	ATGTATGCAC	AAGCACACAG
	TTCGATAGTG TTGTGGTCCC G				
4141	ATGTATGTAC AGCACACACA CA	ACACACACA	CCCCAAAAGG	AGAGAAAAGG	
	TACATACATG TCGTGTGTGT G				TTCTTTTGTA
4201	TTATAAAAAG CGACAGCTAC CO				
	AATATTTTC GCTGTCGATG GC		TTATCAGAAA	<u> </u>	TGTCCTCGAG
4261	TCCATAAGGA ATTATCATGA G			ACTCTCCCAG	
				TGAGAGGGTC	
4321	TGAAGCTGGT CCACRTCTAT A	AACAGGTGA	CACTGGCTGC	AGCAAAAAGC	CATTCGATCC
	ACTTCGACCA GGTGRAGATA T				
4381	ACACAAATTG ATCTTCTATC AT	TCTTGGAAT	CTGAATTGCA	GGGAGGAGCA	GYATGTAAGA
	TGTGTTTAAC TAGAAGATAG TA				
4441	CGACCGTTTA ATTCAGGCAT TO	CCGAAGGCA	TGAGCGCATG	GATTCTRTCA	CCAAGCGTAT
	GCTGGCAAAT TAAGTCCGTA AC				
4501	AAAAGGACCC TGGCATTGGG A	AACCTATGA	CGGACTGTTT	TTGCTGTAGA	AGTAGGGATT
	TTTTCCTGGG ACCGTAACCC TT	TTGGATACT	GCCTGACAAA	AACGACATCT	TCATCCCTAA
4561	TTACAGAAGT CTCCTTGRAT TI	TGCCCTGCC	TGGGGCAGTT	TTGCAGAGGA	ACCTGCCAGA
	AATGTCTTCA GAGGAACRTA AF	ACGGGACGG	ACCCCGTCAA	AACGTCTCCT	TGGACGGTCT
4621	GATTTATTGG CTGGTCAGTC TO	CTTGTGAAA	TAGTATCATG	TGAGAAACAG	TTTGTAGAAA
	CTAAATAACC GACCAGTCAG AC	GAACACTTT	ATCATAGTAC	ACTCTTTGTC	AAACATCTTT
4681	AAAACTATAC CTGGGAAGAC CT	TTTGCAACA	TTGTTCCTTC	CATGGGCCAA	GACTCAGTTA
	TTTTGATATG GACCCTTCTG GA	AAACGTTGT	AACAAGGAAG	GTACCCGGTT	CTGAGTCAAT
4741	GGAGGCATAA ATCTGCCCGG AA	ATAAACTAG	GCCAGGATAC	AGCCATGTTT	AGTTAATAAT
	CCTCCGTATT TAGACGGGCC TT	PATTTGATC	CGGTCCTATG	TCGGTACAAA	TCAATTATTA
• •	EcoRi				
4801	TTCCTTTTAG AATTCACACA GG	CAGGATTG	GTTTTTTGT	GTCTTGGCAA	GTGGAGCATA
	AACCAAAATC TTAAGTGTGT CC	CGTCCTAAC	CAAAAAAACA	CAGAACCGTT	CACCICGIAI
4861	TTTAACATAC AGGCATGGGA AT	CCTGCCTC	TTAGCTTTTC	CCACCCTCTT	GTCTCACCAA
	AAATTGTATG TCCGTACCCT TA	AGGACGGAG	AATCGAAAAG	GGTGGGAGAA	CAGAGIGGII
4921	GTTTTTCTC TCCAAAGGTT TC	CAGGAATT	TCTCATTAAT	GGCTGATGCA	MACTIAGICA MACA BACACA
	CAAAAAAGAG AGGTTTCCAA AG	GTCCTTAA	AGAGTAATTA	CCGACTACG1	*COCY TOTOL
4981	ATAATAATGA ATATAAACAA TG	CTCACCTC	ACCAAAATTA	TATTATTTGC	TCACTABACA
	TATTATTACT TATATTTGTT AC	GAGTGGAG	TGGTTTTAAT	MINNIAMOG.	CTCTCCTTAT
5041	GATAACACAA ATTTTATCGC AA	TGGTTATT	ATTTAATTTG	**CCCCTCTCT	CACACCAATA
	CTATTGTGTT TAAAATAGCG TT	ACCAATAA	TAAATTAAAC	ACCOGIGIGI	ATACCACTGT
5101	CTTTGTTGT GGTTGTTTCT GA	GAAAATGT	TCTTGGATAT	CARTCACCCA	TATEGTCACA
	GAAAACAACA CCAACAAAGA CT	CTTTTACA	AGAACCIAIA	CMITCHOGGI	ATTOTATOTO
5161	GAAGTATTGA TCCCGGGCAG CA	AAATACAG	CCTAAGGTTT	ChTTTCTACT	TAAGATAGAG
	CTTCATAACT AGGGCCCGTC GT	TTTATGTC	CON COCONN N	CTANACTATE	CTGGGCTGGT
5221	AGTTCATCAG AGGGCCTGAG AA	GCTGCGGG	CCAGTGTAAA	CATTACATAC	GACCCGACCA
	AGTTCATCAG AGGGCCTGAG AATTCAAGTAGTC TCCCGGACTC TT	CGACGCCC	CGTCUCUTIT	CCTGTCCCCA	GCTCCCTCCA
5281					
	CCACCAGTCG GAGGGGAACG GT	TOTTOTOT	CONTROCTES.	CATATTCTCC	CATAATGGCA
5341	CGCCTGAAGA GTGACCAGTG CTG GCGGACTTCT CACTGGTCAC GA	GGCCCGAC (CCTACCCIGA	CTATAAGAGG	GTATTACCGT
	AAAAAATAGG CAGTTTGATG TG	* CCBCBCB	A GTGTGGCTC	TCCTCTTTG	AGCATGTGTT
5401	AAAAAATAGG CAGTTTGATG TG	TECRETATI A	CACACCGAG	AGGAGAAAAC	TCGTACACAA
	TTTTTATCC GTCAAACTAC AC	100101111		·	


Figure 15 (con't)

5461	AGCATTTTTA TTTTATACTO	CATCCAGTGAA	CTCTGCTCTT	CCAAGTGTGT	TCATGTATGT
	TCGTAAAAAT AAAATATGAC	TAGGTCACTT	GAGACGAGAA	GGTTCACACA	AGTACATACA
5521	GCTAGATATA TTAGCACAGO	CTGCCTTCTG	CTGCACAACG	CCTTAGAGAC	CCGGCCTTTC
	CGATCTATAT AATCGTGTCC	GACGGAAGAC	GACGTGTTGC	GGAATCTCTG	GGCCGGAAAG
5581	AATGAGCTTA GCTTGTGCTC	TGTTTCTGCT	CTCTTAGGTC	TAAACTATGG	TGTCAGTTTT
	TTACTCGAAT CGAACACGAC	ACAAAGACGA	GAGAATCCAG	ATTTGATACC	ACAGTCAAAA
5641	AATAGAACAA AAGTATGCAT	CTTGCCTTGG	CTTGAGCCTT	TTCGTTTTCA	ATGCTGACTT
	TTATCTTGTT TTCATACGTA	GAACGGAACC	GAACTCGGAA	AAGCAAAAGT	TACGACTGAA
5701	CTCCCCTTTC TCTCCTGTGC	TCACCTTACC	TTTCCAGAGT	GTAAGGGACA	ACTTTTAAG G
	GAGGGGAAAG AGAGGACACG	AGTGGAATGG	AAAGGTCTCA	CATTCCCTGT	TGAAAATTCC
5761	AGGCGTGTCC CTGGTAGGGC	CATCCCTGTT	CACCAGGTGC	CTGTCATCAC	CCCACTTGAC
•	TCCGCACAGG GACCATCCCC	GTAGGGACAA	GTGGTCCACG	GACAGTAGTG	GGGTGAACTG
5821	TGACATCTAC CCTGGTGACT	ATGGGTTCCT	CTTGTTTGTA	GGGAACGGTG	GCTCCAGGTG
	ACTGTAGATG GGACCACTGA	TACCCAAGGA	GAACAAACAT	CCCTTGCCAC	CGAGGTCCAC
5881	CAGCCATCAA TCTGTTGGGT	TCTGGTTCCC	GGCTGCCTTT	GGTTTTGAAA	GTCTCTTCTC
••••	CTCCGTAGTT AGACAACCCA	AGACCAAGGG	CCGACGGAAA	CCAAAACTTT	CAGAGAAGAG
5941	TOTATATATOC TACCOTOCAT	TTGCTTTGTG	TGGTGCTGAT	GCTGTGGCAG	TAGGATCTTG
02.1	ACATATAAGG ATGGGACGTA	AACGAAACAC	ACCACGACTA	CGACACCGTC	ATCCTAGAAC
6001	CATCACTCTC CATCACTCAC	AGACTCCCCC	TGTTGCAAAG	TGTCAGGCTG	ACTCGACAGT
	CTACTGAGAG GTAGTCAGTG	TCTGAGGGGG	ACAACGTTTC	ACAGTCCGAC	TGAGCTGTCA
6061	CACCCEDADA TOTCACTOAC	TCACACACACAG	GCTGTCAGCC	ACGGCTTCCA	CTTGCATGGC
	GTGGCATTTT AGACTCAGTC	AGTGTGTGTC	CGACAGTCGG	TGCCGAAGGT	GAACGTACCG
6121	MANUCEAUTH TOACACCTCA	GTTTCTGTTG	CTGGCTGGCT	GACTGGCATT	ATCTATGCTA
	ATAAGATAAA AGTGTGCACT	CAAAGACAAC	GACCGACCGA	CTGACCGTAA	TAGATACGAT
6181	AGTTGAAATC AGGAGTGTGC	CCAGCAGAGC	CCATCATTCT	CACTGTCTTT	GAAACAAAGC
	TCAACTTTAG TCCTCACACG	GGTCGTCTCG	GGTAGTAAGA	GTGACAGAAA	CTTTGTTTCG
6241	TGTACGGTTT GATCGATGAA	CGTATTTAAA	GCATTTCATG	CAATGACAAA	GTGCTCAGTA
	ACATGCCAAA CTAGCTACTT	GCATAAATTT	CGTAAAGTAC	GTTACTGTTT	CACGAGICAI
6301	GTGGAAGGCA GGCTGTGACC	AGTCTGCCTG	CTCCTTACTA	TAATIGIGAG	CTABACAATC
	CACCTTCCGT CCGACACTGG	TCAGACGGAC	GAGGAATGAT	ATTAACACTC	*CONCA BERT
6361	TGGAACAGTA CATGGAGGCC	TGACCTTGTG	GGGGCACAGG	GTGGAACCTT	TCCDCTTATA
	ACCTTGTCAT GTACCTCCGG	ACTGGAACAC	CCCCGTGTCC	CACCITGGAA	TCGACTIATA
6421	AGTGTGTGTC TCAAGAGGAA	GTCAGGGTAC	TAGCTCAGTG	CTCAATCTCC	TOTAL
	TCACACACAG AGTTCTCCTT	CAGTCCCATG	ATCGAGTCAC	GAGTTAGAGG	CTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT
6481	TATACATTTG CCCGTTTTAT	CTCTAATGTG	AAATAAATCC	CCAAACACTT	CADATAGCAC
	ATATGTAAAC GGGCAAAATA	GAGATTACAC	TTTATTTAGG	GGTTTGTGAA	TEGTCACCCC
6541	TAGCGTACCT AAAAGACTAT	TCTATTATGG	GTGTCCCCAC	TTTCTTGGII	ACCAGTGGGG
	ATCGCATGGA TTTTCTGATA	AGATAATACC	CACAGGGGIG	AAAGAACCAA	
		ledX	- CMC2 CM2 M2	አ አ መር እ ጥር ጥ አ ጥ	GGGAATAGTG
6601	GATCCCCCGG TCTTCTGCTG	TATCTAGAAC	AGTGACTATA	TTACTACATA	CCCTTATCAC
	CTAGGGGCC AGAAGACGAC	ATAGATCTTG	TCACTGATAT	CATTTACTGT	
6661	TTTCCATATG ATCTGTTGTC	TGGAGTATAT	CCATCATGIT	GTAAATGACA	
	AAAGGTATAC TAGACAACAG	ACCTUATATA			
6721	AGTGCAGCTG ATGATGCAAA TCACGTCGAC TACTACGTTT	GCAGTCTCTC	ACACACATOT	CACGGGGTGG	ATAAATTTTT
	TCACGTCGAC TACTACGTTT TCACGTACAA NCCCAGAACA	CGTCAGAGAG	CTTDACATA	GAAACAAACG	CAGCGTCTGG
6781	TCACGTACAA NCCCAGAACA AGTGCATGTT NGGGTCTTGT	CTGTGAAACA	CAATTGTATT	CTTTGTTTGC	GTCGCAGACC
	AGTGCATGTT NGGGTCTTGT	GUCUCITIGI	<u></u>		

Figure 15 (con't)

6841	ATTCTTTCCA	AGGAGAGCAG	CTTTCTCCAC	AGGAACACAG	TAACAAAAGA	GGTCCGCCGC
	TAAGAAAGGT	TCCTCTCGTC	GAAAGAGGTG	TCCTTGTGTC	ATTGTTTTCT	CCAGGCGGCG
6901	CATCCACACC	CAGCCAAGAC	ACCTCAGAGG	CCATAGGGAC	AACCTCCTTG	CTGGCCAACA
	GTAGGTGTGG	GTCGGTTCTG	TGGAGTCTCC	GGTATCCCTG	TTGGAGGAAC	GACCGGTTGT
6961	CCTGCTGGAG	CAGGGCACAG	GTCCCAGCAA	CTGATCCTCA	GTGGATGGGT	CCGCAGTCAA
	GGACGACCTC	GTCCCGTGTC	CAGGGTCGTT	GACTAGGAGT	CACCTACCCA	GGCGTCAGTT
	***************************************				Hindill	EcoRY
7021	AGCCTTAATG	GGCTCTCTTT	TGAAGGGGAA	AGAAANNTTT		
	TCGGAATTAC	CCGAGAGAAA	ACTTCCCCTT	TCTTTNNAAA	GTTCGAATAC	TATAGGTTGT
7081	TTATTATAGT	TGATGAGTTA	GTAAATTCCG	AAAAAAAAG	ATGATTTTAT	ATGTATGACA
	AATAATATCA	ACTACTCAAT	CATTTAAGGC	TTTTTTTTC	TACTAAAATA	TACATACTGT
7141	TAAAAAAAT	CTTTGTAAAG	TGCGCAAGTG	CAATAATTTA	AAGAGGTCTT	ATCTTTGCAT
	ATTTTTTTA	GAAACATTTC	ACGCGTTCAC	GTTATTAAAT	TTCTCCAGAA	TAGAAACGTA
7201	TTATAAATTA	TAAATATTGT	ACATGTGTGT	AATTŢTTCAT	GTATTCATTT	GCAGTCTTTG
	AATATTTAAT	ATTTATAACA	TGTACACACA	TTAAAAAGTA	CATAAGTAAA	CGTCAGAAAC
7261	TATTTAAAAA	AACTTTACTG	TTATGTTTGT	ATAATAGAAC	ATTAATCATT	TATTATAACT
	ATAAATTTT	TTGAAATGAC	AATACAAACA	TATTATCTTG	TAATTAGTAA	ATAATATTGA
7321	CAGACAAGGT	GTAAATAAAT	TCATAATTCA	AACAGCCAGT	ATATATGCAT	ATATGGGTGT
	GTCTGTTCCA	CATTTATTTA	AGTATTAAGT	TTGTCGGTCA	TATATACGTA	TATACCCACA
7381	TACATTGCAA	AAATCTCTAT	CTTTGTTCTA	TTCACATGCT	TAAAGAAGTA	AGAAATCTTT
	ATGTAACGTT	TTTAGAGATA	GAAACAAGAT	AAGTGTACGA	ATTTCTTCAT	TCTTTAGAAA
7441	TGTGGATATG	TAATTATACA		ATATATATGT		
	ACACCTATAC	ATTAATATGT	ATATTTCATA	TATATATACA	TACTATGTAC	TTTATATAAA
7501		AATTTTAA	200:::::	TTGGTGTGAA		
	TCTTTACAAG	TATTAAAATT	ACCTATAAGA	AACCACACTT .	ATTAACTTAT	GTTGTAAAAA
7561	AAAATGAAAA	AAAAAAAA	С			
	TTTTACTTTT	TTTTTTTTT	G [.]			


Figure 16



PDE10a and RACEs compiled 7581 bp


Figure 17

PDE10A compiled - coding sequence and features



34/41

Figure 18
PDE10A compiled - restriction sites



35/41

Figure 19

1						CTGAGCTGGA
				CAGTTGGACT	•	
61				TCCAGGAAGA		
				AGGTCCTTCT	• •	
121				TCTGTATACT		
				AGACATATGA		
181				CGCGCCCTGA		
				GCGCGGGACT		
241				TGGTTTGACG		
				ACCAAACTGC	*** ** *** ***	
301				ATTTGTTTCT		
				TAAACAAAGA		
361				CAAAGCAAAA		
				GTTTCGTTTT		
421				GGGAGTCGTG		
				CCCTCAGCAC		
481	AGAGCAGCGC	CTGGACACGG	GCGGGGACAA	CCACCTGCTC	CTCTATGAGC	TCAGCAGCAT "
				GGTGGACGAG		
541	CATCAGGATA	GCCACAAAAG	CCGACGGATT	TGCACTGTAC	TTCCTTGGAG	AGTGCAATAA
				ACGTGACATG		
601	TAGCCTGTGT	GTGTTCATAC	CACCCGGGAT	GAAGGAAGGC	CAACCCCGGC	TCATCCCTGC
				CTTCCTTCCG		
661	AGGGCCCATC	ACCCAGGGTA	CCACCATCTC	TGCCTACGTG	GCCAAGTCTA	GGAAGACGTT
				ACGGATGCAC		
721	GTTGGTAGAG	GATATCCTTG	GGGATGAGCG	ATTTCCTCGA	GGTACTGGCC	TGGAATCAGG
				TAAAGGAGCT		
781	AACCCGCATC	CAGTCTGTTC	TTTGCTTGCC	CATTGTCACT	GCCATTGGAG	ACTTGATTGG
				GTAACAGTGA		
841	CATCCTTGAA	CTGTACAGGC	ACTGGGGCAA	AGAGGCCTTC	TGCCTCAGCC	ATCAGGAGGT
				TCTCCGGAAG		
901	TGCAACAGCC	AATCTTGCTT	GGGCTTCCGT	AGCAATACAC	CAGGTGCAGG	TGTGTAGAGG
				TCGTTATGTG		
961	TCTCGCCAAA	CAGACCGAAC	TGAATGACTT	CCTACTCGAC	GTATCAAAGA	CATACTTTGA
				GGATGAGCTG		
1021	TAACATAGTT	GCCATAGACT	CTCTACTTGA	ACACATCATG	ATATATGCAA	AAAATCTAGT.
				TGTGTAGTAC		
1081	GAACGCCGAC	CGCTGCGCGC	TCTTCCAGGT	GGACCACAAG	AACAAGGAGC	TGTACTCGGA
	CTTGCGGCTG	GCGACGCGCG	AGAAGGTCCA	CCTGGTGTTC	TTGTTCCTCG	ACATGAGCCT
1141	CCTGTTTGAC	ATTGGGGAGG	AGAAGGAGGG	GAAGCCCATC	TTCAAGAAGA	CCAAGGAGAT
				CTTCGGGTAG		
1201	CAGATTTTCC	ATTGAGAAAG	GGATTGCTGG	TCAAGTGGCA	AGAACAGGCG	AAGTCTTGAA
	GTCTAAAAGG	TAACTCTTTC	CCTAACGACC	AGTTCACCGT	TCTTGTCCGC	TTCAGAACTT
1261	CATTCCCGAT	GCCTACGCGG	ACCCTCGCTT	TAACAGGGAG	GTGGACCTGT	ACACAGGCTA
	GTAAGGGCTA	CGGATGCGCC	TGGGAGCGAA	ATTGTCCCTC	CACCTGGACA	TGTGTCCGAT
1321	CACCACGAGG	AACATTCTGT	GTATGCCCAT	AGTGAGCCGA	GGCAGCGTGA	TTGGCGTGGT
	GTGGTGCTCC	TTGTAAGACA	CATACGGGTA	TCACTCGGCT	CCGTCGCACT	AACCGCACCA

Figure 19 (con't)

1381	GCAGATGGTG AACAAGATCA	GCGGTAGCGC	CTTCTCCAAG	ACAGACGAGA	ACAACTTCAA
1301	CGTCTACCAC TTGTTCTAGT	CGCCATCGCG	GAAGAGGTTC	TGTCTGCTCT	TGTTGAAGTT
1441	GATGTTTGCT GTCTTCTGCG	CACTGGCCTT	GCACTGTGCT	AACATGTACC	ACAGGATCCG
1.11	CTACAAACGA CAGAAGACGC	GTGACCGGAA	CGTGACACGA	TTGTACATGG	TGTCCTAGGC
1501	CCACTCAGAA TGCATCTACA	GGGTTACCAT	GGAGAAGCTT	TCCTACCACA	GCATCTGCAC
1001	GGTGAGTCTT ACGTAGATGT	CCCAATGGTA	CCTCTTCGAA	AGGATGGTGT	CGTAGACGTG
1561	CTCCGAGGAG TGGCAAGGCC	TCATGCGCTT	CAACCTACCA	GCACGCATCT	GCCGGGACAT
2002	GAGGCTCCTC ACCGTTCCGG	AGTACGCGAA	GTTGGATGGT	CGTGCGTAGA	CGGCCCTGTA
1621	CGAGCTATTC CACTTTGACA	TTGGTCCTTT	CGAGAACATG	TGGCCTGGGA	TCTTTGTCTA
	GCTCGATAAG GTGAAACTGT	AACCAGGAAA	GCTCTTGTAC	ACCGGACCCT	AGAAACAGAT
1681	CATGATCCAT CGGTCTTGTG	GGACATCCTG	TTTTGAACTT	GAAAAATTGT	GCCGTTTTAT
	GTACTAGGTA GCCAGAACAC	CCTGTAGGAC	AAAACTTGAA	CTTTTTAACA	CGGCAAAATA
1741	CATGTCTGTG AAGAAGAACT	ATCGGCGGGT	TCCTTACCAC	AACTGGAAGC	ATGCAGTCAC
	GTACAGACAC TTCTTCTTGA				
1801	GGTGGCACAC TGCATGTATG	CCATACTTCA	AAACAACAAT	GGCCTCTTCA	CAGACCTCGA
	CCACCGTGTG ACGTACATAC	GGTATGAAGT	TTTGTTGTTA	CCGGAGAAGT	GTCTGGAGCT
1861	GCGCAAAGGC CTGCTAATTG	CGTGTCTGTG	CCATGACCTG	GACCACAGGG	GCTTCAGTAA
	CGCGTTTCCG GACGATTAAC	GCACAGACAC	GGTACTGGAC	CTGGTGTCCC	CGAAGTCATT
1921	CAGCTACCTG CAGAAGTTCG	ACCACCCCT	GGCGGCGCTG	TACTCCACCT	CCACCATGGA
	GTCGATGGAC GTCTTCAAGC	TGGTGGGGGA	CCGCCGCGAC	ATGAGGTGGA	GGTGGTACCT
1981	GCAACACCAC TTCTCCCAGA	CGGTGTCCAT	CCTTCAGCTG	GAAGGGCACA	ATATCTTCTC
	CGTTGTGGTG AAGAGGGTCT	GCCACAGGTA	GGAAGTCGAC	CTTCCCGTGT	TATAGAAGAG
2041	CACCTGAGC TCCAGCGAGT	ACGAGCAGGT	GCTGGAGATC	ATCCGCAAAG	CCATCATCGC
	GTGGGACTCG AGGTCGCTCA	TGCTCGTCCA	CGACCTCTAG	TAGGCGTTTC	GGTAGTAGCG
2101	CACCGACCTC GCCCTATACT	TTGGGAACAG	GAAGCAGTTG	GAGGAGATGT	ACCAGACAGG
	GTGGCTGGAG CGGGATATGA	AACCCTTGTC	CTTCGTCAAC	CTCCTCTACA	TGGTCTGTCC
2161	GTCGCTGAAC CTCCACAACC	AGTCCCATCG	AGACCGTGTC	ATCGGCTTGA	TGATGACTGC
	CAGCGACTTG GAGGTGTTGG	TCAGGGTAGC	TCTGGCACAG	TAGCCGAACT	ACTACTGACG
2221	CTGTGATCTT TGCTCTGTGA	CCAAACTATG	GCCAGTTACA	AAATTGACAG	CCTTACTATA
	GACACTAGAA ACGAGACACT	GGTTTGATAC	CGGTCAATGT	TTTAACTGIC	BCCCCATTCC.
2281	ATATGCAGAA TTCTGGGCTG	AGGGTGATGA	GATGAAGAAG	CACCCCTATC	TCGGGTAAGG
	TATACGTCTT AAGACCCGAC	TCCCACTACT	CTACTTCTTC	GACCCGIAIG	TCGGGTTMGC
2341	TATGATGGAC AGAGACAAGC	GAGATGAAGT	CCCTCAAGGG	CRCCACCCTA	AGATGTTACG
	ATACTACCTG TCTCTGTTCG	CTCTACTTCA	GGAGTICCC	CONCOCNONC	ACCOTOTOT
2401	TGTGGCCATT CCCTGCTATA ACACCGGTAA GGGACGATAT	CCACCTTGAC	CCTCTACCAC	CCACCCACAG	TCGGAGACGA
	ACACCGGTAA GGGACGATAI	GGTGGAACTG	CGICIAGGAG	ATTCCCCCC	AAGAGACAGC
2461	GAAGGCCTGC AGGGATAACC	TCAATCAGIG	CCTCTTCCAT	TAAGCGCCCC	TTCTCTGTCG
	CTTCCGGACG TCCCTATTGG AATGTGGATT TCAGGCCCAG	AGTIAGICAC	TACCAACACC	ACACCTGAGA	AGCTGAACGT
2521	TTACACCTAA AGTCCGGGTC	CCCCCCCCC	ATCGTTCTCG	TGTGGACTCT	TCGACTTGCA
	GAAGGTTGAA GACTGATCCT	CANCECACCE	ССТСАТСТСТ	GCCCAGCAAC	CGACTCAACC
2581	GAAGGTTGAA GACTGATCCT CTTCCAACTT CTGACTAGGA	CTTCACTGCA	GGACTACAGA	CGGGTCGTTG	GCTGAGTTGG
	TGCTTCTGTG ACTTCGTTCT		AAGGGGTGAA	AACCCCCTGT	CAGAAGGTAC
2641	ACGAAGACAC TGAAGCAAGA	AAAACAAAAG	TTCCCCACTT	TTGGGGGACA	GTCTTCCATG
0701	CGTCGCATAT CCATGTGAAG	CAGACGACTC	CCTGCTTGCC	GCACACACCT	CGGACAGTGA
2701	GCAGCGTATA GGTACACTTC	GTCTGCTGAG	GGACGAACGG	CGTGTGTGGA	GCCTGTCACT
	GCAGCGIIII. GGII.GII		- , · · · · · · · ·	•	

Figure 19 (con't)

2761	GCAACCCAG	CTCTGCCGTG	TTCAGACGTC	GGCTACTCCG	TGGCTCCACC	TGACCTCCGA
	CGTTGGGTC	GAGACGGCAC	AAGTCTGCAG	CCGATGAGGC	ACCGAGGTGG	ACTGGAGGCT
2821				TGTCTGGAGG		
				ACAGACCTCC	·- ·- · · · ·	
2881				GGCCAGTTCC		
		· · · · · · · · · · · · · · · · · · ·		CCGGTCAAGG		
2941				CACGCCCCTT		
			···· · · · ·	GTGCGGGGAA		
3001			•	TGGGACACAT		
				ACCCTGTGTA		
3061				ACAGGTGCAC		
				TGTCCACGTG		·
3121				CACTCAGAAG		
				GTGAGTCTTC		
3181				CGGGGAACTG		
2041				GCCCCTTGAC	· · · · - · · · · · ·	
3241				TCACTGTAGC		
2201				AGTGACATCG		
3301				TATAAATTTC ATATTTAAAG		
3361				ACCTCTATAT		
3301				TGGAGATATA		
3421		·		ATTCCAGAAG		
3421				TAAGGTCTTC		
3481		· · · · · · · · · · · · · · · · · · ·		GGTTCTGATC		
				CCAAGACTAG		
3541	AGGCAGGACA	GGAGGAGGTG	CAGGGCTACC	ACATTGACCC	AGATGGTATC	TCCTCTCACC
				TGTAACTGGG		
3601	ATTCAGACAT	CCATAAGGAA	TGCCAAATGC	TGTATTGAAT	AGTTCTCCTG	TGTGACTTTC
	TAAGTCTGTA	GGTATTCCTT	ACGGTTTACG	ACATAACTTA	TCAAGAGGAC	ACACTGAAAG
3661	TAGAGAAGCC	AGGACACCCC	TGAGCCTTTC	CTGGGAACTC	CTAAGGAAGT	CACAGGTTCA
	ATCTCTTCGG	TCCTGTGGGG	ACTCGGAAAG	GACCCTTGAG	GATTCCTTCA	GTGTCCAAGT
3721	CACCGTGGGG	ATTTTCAGGA	TAGCATGGAG	ACCAGAGAAT	CCCGGTTCGG	TTGTTCTCAC
	GTGGCACCCC	TAAAAGTCCT	ATCGTACCTC	TGGTCTCTTA	GGGCCAAGCC	AACAAGAGTG
3781	TCGGTGAGCC	TTGAGAAGGA	AGAGACTGAC	CAGAAACACT	CACTCAGCAC	TCTGGCAGGA
	AGCCACTCGG	AACTCTTCCT	TCTCTGACTG	GTCTTTGTGA	GTGAGTCGTG	AGACCGTCCT
3841				GGGATAGATT		
				CCCTATCTAA		
3901				TTCAGTTTCC		
		····		AAGTCAAAGG		
3961				CTGTACATGC		
4001	- · 			GACATGTACG		
4021				CTACTCCATT		
4003				GATGAGGTAA		
4081				TGTGCACACG ACACGTGTGC		
······································	GIIGIICICI	.commoigi	191001000	Povedigiec.	Wedferer .	NOW TWO GIGI

Figure 19 (con't)

	4141	AGCACACAGA	TGTATGTACA	GCACACACAC	ACACACACAC	CCCAAAAGGA	GAGAAAAGGA
		TCGTGTGTCT	ACATACATGT	CGTGTGTGTG	TGTGTGTGTG	GGGTTTTCCT	CTCTTTTCCT
	4201			GACAGCTACC			
		TCTTTTGTAA	ATATTTTTCG	CTGTCGATGG	GGGTATAAGT	TTTTATCAAG	AAAAGGGACA
	4261	AGGGAAACAG	GTAGCTCTCC	ATAAGGAAAT	TATCATGAGT	GTGTTCTCCC	ATCAGTGCAC
		TCCCTTTGTC	CATCGAGAGG	TATTCCTTTA	ATAGTACTCA	CACAAGAGGG	TAGTCACGTG
	4321			GAAGCTGGTC			
		AAGAGGGTCC	CCACGAGTGA	CTTCGACCAG	GTGCAGATAT	TTGTCCACTG	TGACCGACGT
	4381			CACAAATTGA			
	.	CGTTTTTCGG	TAAGCTAGGT	GTGTTTAACT	AGAAGATAGT	AGAACCTTAG	ACTTAACGTC
	4441			GACCGTTTAA			
	· ·	CCTCCTCGTC	GTACATTCTG	CTGGCAAATT	AAGTCCGTAA	GGCTTCCGTA	CTCGCGTACC
	4501			AAAGGACCCT			
			·	TTTCCTGGGA			
	4561			TACAGAAGTC			
		ACGACATCTT	CATCCCTAAA	ATGTCTTCAG	AGGAACCTAA	ACGGGACGGA	CCCCGTCAAA
	4621			ATTTATTGGC			
	••·· • · • • ·			TAAATAACCG			
	4681			AAACTATACC			
				TTTGATATGG	 ~~~~~~~~~ ~ ·	·	
	4741			GAGGCATAAA			
				CTCCGTATTT			
	4801			$\mathbf{T}\mathbf{G}\mathbf{G}\mathbf{T}\mathbf{T}\mathbf{T}\mathbf{A}\mathbf{G}\mathbf{A}$			
				ACCAAAATCT			
	4861	TCTTGGCAAG	TGGAGCATAT	TTAACATACA	GGCATGGGAA	TCCTGCCTCT	TAGCTTTTCC
				AATTGTATGT			
	4921			TTTTTTCTCT			
				AAAAAAGAGA			
	4981	GCTGATGCAA	ACTTAGTGAA	TAATAATGAA	TATAAACAAT	CCACCTCA	CCAAAATTAT
				ATTATTACTT			
	5041	ATTATTTGCA	GTCATTTGTG	ATAACACAAA	TTTTATCGCA	TACCADANA	AAATTAAACA
-							CTTGGATATG
	5101			AAAACAACAC			
-							
	5161	TAAGTGCCAA	AUCCAGIGIG	AAGTATTGAT TTCATAAÇTA	CCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	TTTTATCTCC	GATTCCAAAC
				GTTCATCAGA			
	5221	TAAACATCAA	AACATOTCA	CAAGTAGTCT	CCCGGACTCT	TCGACGCCCC	GTCACATTTC
-	6001	MANA CONTOCC	MCCCCOMCCMC	GTGGTCAGCC	#CCCCTTGCC	AAGAAGAGAG	CAATTGAATC
	5281	ATTTCATACG	ACCCGACCAC	CACCAGTCGG	AGGGGAACGG	TTCTTCTCTC	GTTAACTTAG
	5341			GCCTGAAGAG			
	224T	GACAGGGGTC	GAGGGAGGTG	CGGACTTCTC	ACTGGTCACG	ACCGGGCTGC	CTAGCGACTC
	5401			AAAAATAGGC			
		TATAAGAGGG	TATTACCGTT	TTTTTATCCG	TCAAACTACA	CTGGACAAAT	CACACCGAGA
٠	5461			GCATTTTAT			
		GGAGAAAACT	CGTACACAAT	CGTAAAAATA	AAATATGAGT	AGGTCACTTG	AGACGAGAAG

Figure 19 (con't)

					•	
5521	CAAGTGTGT	T CATGTATGTO	CTAGATATAT	TAGCACAGCC	TGCCTTCTGC	TGCACAACGC
			GATCTATATA			
5581	CTTAGAGAC	C CGGCCTTTCA	ATGAGCTTAG	CTTGTGCTCT	GTTTCTGCTC	TCTTAGGTCT
			TACTCGAATC			
5641			ATAGAACAAA			
			TATCTTGTTT	- · · · · · ·		
5701			TCCCCTTTCT			
			AGGGGAAAGA		<u>-</u>	
5761			GGCGTGTCCC			
			CCGCACAGGG	· ··-· · · · · · · · · · · · · · · ·	• •-	
5821			GACATCTACC			
			CTGTAGATGG	·		
5881			AGGCATCAAT			
	·		TCCGTAGTTA	· · · · · · · · · · ·		
5941			GTATATTCCT			
		× - · — · — — — — — — — — — — — — — — — —	CATATAAGGA			
6001			ATGACTCTCC			
			TACTGAGAGG			***************************************
6061			ACCGTAAAAT			
			TGGCATTTTA			• • • • • • • • • • • • • • • • • • • •
6121			ATTCTATTTT			
			TAAGATAAAA		• • •• • •	******
6181			GTTGAAATCA			
			CAACTTTAGT		.,	
6241			GTACGGTTTG			
			CATGCCAAAC		· · · ·	
6301			TGGAAGGCAG			
			ACCTTCCGTC			
6361			GGAACAGTAC			
			CCTTGTCATG			• • • • • • • • • • • • • • • • • • • •
6421			GTGTGTGTCT		•	
			CACACACAGA	· · _		
6481			ATACATTTGC			
			TATGTAAACG		·· · · - ·	
6541			AGCGTACCTA			
		···	TCGCATGGAT	•· • •··• •· • · • · • · •		· · · · · ·
6601			ATCCCCCGGT			
			TAGGGGGCCA		•	
6661			TTCCATATGA			
	TACTACATAC	CCTTATCACA	AAGGTATACT	AGACAACAGA	CCTCATATAC	GATGTACAAG
6721			GTGCAGCTGA			
	·		CACGTCGACT		• • • • • • • • • • • • • • • • • • • •	
6781	•		CACGTACAAN			
			GTGCATGTTN			
6841			TTCTTTCCAA			
	TTTGTTTGCG	TCGCAGACCT	AAGAAAGGTT	CCTCTCGTCG	AAAGAGGTGT	CCTTGTGTCA

Figure 19 (con't)

6901	AACAAAAGAG GTCCGCCGC	C ATCCACACCC	AGCCAAGACA	CCTCAGAGGC	CATAGGGACA
	TTGTTTTCTC CAGGCGCG	TAGGTGTGGG	TCGGTTCTGT	GGAGTCTCCG	GTATCCCTGT
6961	ACCTCCTTGC TGGCCAACA	CTGCTGGAGC	AGGGCACAGG	TCCCAGCAAC	TGATCCTCAG
	TGGAGGAACG ACCGGTTGT	GACGACCTCG	TCCCGTGTCC	AGGGTCGTTG	ACTAGGAGTC
7021	TGGATGGGTC CGCAGTCAA	A GCCTTAATGG	GCTCTCTTTT	GAAGGGGAAA	GAAANNTTTC
*** ,, ,,	ACCTACCCAG GCGTCAGTT	CGGAATTACC	CGAGAGAAAA	CTTCCCCTTT	CTTTNNAAAG
7081	AAGCTTATGA TATCCAACA	TATTATAGTT	GATGAGTTAG	TAAATTCCGA	AAAAAAAAGA
	TTCGAATACT ATAGGTTGT	ATAATATCAA	CTACTCAATC	ATTTAAGGCT	TTTTTTTTCT
7141	TGATTTTATA TGTATGACA	T AAAAAAATC	TTTGTAAAGT	GCGCAAGTGC	AATAATTAA
	ACTAAAATAT ACATACTGTA	TTTTTTTAG	AAACATTTCA	CGCGTTCACG	TTATTAAATT
7201	AGAGGTCTTA TCTTTGCATT	TATAAATTAT	AAATATTGTA	CATGTGTGTA	ATTTTTCATG
	TCTCCAGAAT AGAAACGTAA	ATATŤTAATA	TTTATAACAT	GTACACACAT	TAAAAAGTAC
7261	TATTCATTTG CAGTCTTTGT	ATTTAAAAAA	ACTTTACTGT	TATGTTTGTA	TAATAGAACA
	ATAAGTAAAC GTCAGAAACA	TAAATTTTTT	TGAAATGACA	ATACAAACAT	ATTATCTTGT
7321	TTAATCATTT ATTATAACTO	AGACAAGGTG	TAAATAAATT	CATAATTCAA	ACAGCCAGTA
	AATTAGTAAA TAATATTGAG	TCTGTTCCAC	ATTTATTTAA	GTATTAAGTT	TGTCGGTCAT
7381	TATATGCATA TATGGGTGTT	ACATTGCAAA	AATCTCTATC	TTTGTTCTAT	TCACATGCTT
	ATATACGTAT ATACCCACAA	TGTAACGTTT	TTAGAGATAG	AAACAAGATA	AGTGTACGAA
·7441 ·	AAAGAAGTAA GAAATCTTTI	GTGGATATGT	AATTATACAT	ATAAAGTATA	TATATATGTA
	TTTCTTCATT CTTTAGAAAA	CACCTATACA	TTAATATGTA	TATTTCATAT	ATATATACAT
7501	TGATACATGA AATATATTA	GAAATGTTCA	TAATTTTAAT	GGATATTCTT	TGGTGTGAAT
	ACTATGTACT TTATATAAAT	CTTTACAAGT	ATTAAAATTA	CCTATAAGAA	ACCACACTTA
7561	AATTGAATAC AACATTTTTA	AAATGAAAAA	АААААААА	АААААААА	AAAAAAA
<u></u>	TTAACTTATG TTGTAAAAAT	TTTACTTTTT	TTTTTTTTT	TTTTTTTTT	TTTTTTT

SEQUENCE LISTING

<110> ROBERTSON, Harold
 DENOVAN-WRIGHT, Eileen
 NOVANEURON, INC.

<120> GENE NECESSARY FOR STRIATAL FUNCTION, USES THEREOF, AND COMPOUNDS FOR MODULATING SAME

<130> 36541-0005

<140>

<141>

<150> US60/158,043

<151> 1999-10-07

<150> US60/217,765

<151> 2000-07-12

<160> 12

<170> PatentIn Ver. 2.0

<210> 1

<211> 3236

<212> DNA

<213> mouse

<400> 1

cactgaaget ggtccacgte tataaacagg tgacactgge tgeageaaaa agecattega 60
tecacacaaa ttgatettet ateatettgg aatetgaatt geaggagga geagtatgta 120
agaegaeegt ttaatteagg catteegaag geatgagege atggattetg teaceaageg 180
tataaaagga eeetggeatt gggaaaeeta tgaeggaetg tttttgetgt agaagtaggg 240
attttacaga agteteettg aatttgeeet geetggggea gttttgeaga ggaacetgee 300
agagatttat tggetggtea gtetettgtg aaatagtate atgtgagaaa eagtttgtag 360
aaaaaaaeta tacetgggaa gaeetttgea aeattgttee tteeatggge caagaeteag 420
ttaggaggea taaatetgee eggaataaae taggeeagga tacageeatg tttagttaat 480

aatttggttt tagaattcac acaggcagga ttggtttttt tgtgtcttgg caagtggagc 540 atatttaaca tacaggcatg ggaatcetge etettagett tteecaceet ettgteteac 600 caagtttttt ctctccaaag gtttccagga atttctcatt aatggctgat gcaaacttag 660 tgaataataa tgaatataaa caatgctcac ctcaccaaaa ttatattatt tgcagtcatt 720 tgtgataaca caaattttat cgcaatggtt attatttaat ttgtggccac acactgtggt 780 tatcttttgt tgtggttgtt tctgagaaaa tgttcttgga tatgtaagtg ccaataccag 840 tgtgaagtat tgatcccggg cagcaaaata cagcctaagg tttgtaaaca tcaattctat 900 ggtggtggtc agcctcccgc ctgaagagtg accagtgctg gcccgacgga tcgctgagat 1020 atteteceat aatggeaaaa aaataggeag tttgatgtga eetgtttagt gtggetetee 1080 tettttgage atgtgttage atttttattt tataeteate eagtgaaete tgetetteea 1140 agtgtgttca tgtatgtgct agatatatta gcacagcctg ccttctgctg cacaacgcct 1200 tagagacccg gcctttcaat gagcttagct tgtgctctgt ttctgctctc ttaggtctaa 1260 actatggtgt cagttttaat agaacaaaag tatgcatctt gccttggctt gagccttttc 1320 gttttcaatg ctgacttctc ccctttctct cctgtgctca ccttaccttt ccagagtgta 1380 agggacaact tttaaggagg cgtgtccctg gtaggggcat ccctgttcac caggtgcctg 1440 teateacece aettgaetga catetacect ggtgaetatg ggtteetett gtttgtaggg 1500 aacggtggct ccaggtggag gcatcaatct gttgggttct ggttcccggc tgcctttggt 1560 tttgaaagtc tcttctctgt atattcctac cctgcatttg ctttgtgtgg tgctgatgct 1620 gtgcgcagta ggattcttgg atgactctcc atcagtcaca gactcccct gttgcaaagt 1680 gtcaggctga ctcgacagtc accgtaaaat ctgagtcagt cacacacagg ctgtcagcca 1740 eggetteeae ttgeatgget attetatttt cacaegtgag tttetgttge tggetggetg 1800 actggcatta tctatgctaa gttgaaatca ggagtgccca gcagagccca tcattctcac 1860 tgtctttgaa acaaagctgt acggtttgat cgatgaacgt atttaaagca tttcatgcaa 1920 tgacaaagtg ctcagtagtg gaaggcaggc tgtgaccagt ctgcctgctc cttactataa 1980 ttgtgaggat ttgttactgg aacagtacat ggaggcctga ccttgtgggg gcacagggtg 2040 gaaccttagc tgaatatagt gtgtgtctca agaggaagtc agggtactag ctcagtgctc 2100 aatctccagg tactatatat acatttgccc gttttatctc taatgtgaaa taaatcccca 2160 aacacttgtt tatcgtgtag cgtacctaaa agactattct attatgggtg tccccacttt 2220 cttggtttgg tcaccccgat cccccggtct tctgctgtat ctagaacagt gactataaat 2280 gatgtatggg aatagtgttt ccatatgatc tgttgtctgg agtatatgct acatgttcaa 2340 ttactgtaca aaaacccagt gcagctgatg atgcaaagca gtctctctct gtgtacagtg 2400 ccccacctat ttaaaaatca cgtacaascc cagaacactg tgaaacactt aacataagaa 2460 caaacgcagc gtctggattc tttccaagga gagcagcttt ctccacagga acacagtaac 2520 aaaagaggtc cgccgccatc cacacccagc caagacacct cagaggccat agggacaacc 2580 teettgetgg ccaacacetg etggageagg ggeacaggte ccageaactg atceteagtg 2640 gatgggtccg cagtcaaagc cttaatgggc tctcttttga aggggaaaga aagaatttca 2700 agcttatgat atccaacatt attatagttg atgagttagt aaattccaaa aaaaaaagat 2760 gattttatat gtatgacata aaaaaaatct ttgtaaagtg cgcaagtgca ataatttaaa 2820

<210> 2

<211> 5752

<212> DNA

<213> mouse

<400> 2

aagtgtaaat aaaataaaca tctaataaaa aaaattacat accatagagg aacaagataa 60 tttctgccca acttcatacc ctccagcgta tagtgttgag gtttggtctg ttgctgtgta 120 ttgtaatgta atgttaaatt ctctacctga aggtctaggc ctacaagtga attctcatgt 180 ttatagagtt ttgttgtgca aaccttgttc cttaatttaa aactatggtt aaaaaacaaa 240 acaaaactgg ctacagccaa taactgaagg gggttacctt gttgaagggg tggaaaagag 300 acgaggggag atgggccatg agaacttggc caggagaaat agccagtatc tggagtacac 420 cactgaggag gtagccaggc tagcagttag aagagtagat taggggttat ttttccccca 480 ctccacatag ttatcaaagc caaataaaat aaccatagtc tgagtctcat ctatttgtaa 540 gctagttggg tataagatta atttggctgt actacagttt agatttctaa cataggaact 600 atcaaaaact tgctcaaaca agaacatgct gacaatattt taaaatgatt atttatattg 660 tttgcacttt ctaaagtttc ttctaaatgt tccatggtca aattaaaaaa tatacatatt 720 ggctattaaa ttcgtctaag tggggctgga gagatagctc agaggttaag agcactgact 780 gctcttccag aggtcctgag ttcaattccc agcgaccaca tggtggctca cagccatctg 840 taatagatag gatctgacgc cctcttctgg agtgtctgaa gacagctaca atgtactcat 900 atatattaaa taaataatat tagaaaatto ttotaagtgt atcatttata gaatatttaa 960 tatataaagt aaatgcctca ggaaatataa acttggaatt aaatcaaaga acttcatgag 1020° tagtgggcca caaaaaatgt gtaccagggg aagaccggag ggaggggaga aggaagggat 1080 ggagatagaa ttttgcctct gcattccttg ggctggcaca ggtataatgc tgtgggaatt 1140 gggaaactac aaggaagctg caaagctggg cggaactcgt ttccgcaagc tgggctcatc 1200 taagtgtcca tgcatggctg ccacactgca gtgaacttta aaacatttgt gttccagaga 1260 tgtagagatg ctcacaatag tacaaaggcg ggagggaggt atttccagac taagaggaag 1320 aaaaaccatt gctgattaaa catctgcata tgagcgcccc cacctccata cacacaca 1380 cacacacaca cacacaca caaccaaaca gaacaaatac acatgcatgt ctacagcctg 1440 caggaacaaa atggtatgtc tgtgaggaac caggagatgc acaggtccta acctctgtct 1500

cctacaagcc ctgaagtctg gtcagggtca aatgtacaaa agcaggctaa ggaagctgtt 1560 tagtgaaaga tttttttctt caactctagg aacaacctat ttcctaggat ttggagagtg 1620 ctcaggagga aacattcaga caactgatgc tctctgtgta ccccagattc aggtattggg 1680 gtagttagtt gtgctcatgt atgtgctaga tatattagca cagcctgcct tctgctgcac 1740 aacgeettag agacceggee ttteaatgag ettagettgt getetgttte tgetetetta 1800 ggtctaaact atggtgtcag ttttaataga acaaaagtat gcatcttgcc ttggcttgag 1860 cettttegtt tteaatgetg aetteteece ttteteteet gtgeteaeet taeettteea 1920 gagtgtaagg gacaactttt aaggaggegt gteeetggta ggggeateee tgtteaceag 1980 gtgcctgtca tcaccccact tgactgacat ctaccctggt gactatgggt tcctcttgtt 2040 tgtagggaac ggtggctcca ggtggaggca tcaatctgtt gggttctggt tcccggctgc 2100 ctttggtttt gaaagtetet tetetgtata tteetaceet geatttgett tgtgtggtge 2160 tgatgctgtg cgcagcagga ttcttggatg actctccatc agtcacagac tccccctgtt 2220 gcaaagtgtc aggctgactc gacagtcacc gtaaaatctg agtcagtcac acacaggctg 2280 teageracgg ctteracttg catggetatt ctatttteac acgtgagttt ctgttgctgg 2340 ctggctgact ggcattatct atgctaagtt gaaatcaggg gtgcccagca gagcccatca 2400 ttctcactgt ctttgaaaca aagctgtacg gtttgatcga tgaacgtatt taaagcattt 2460 catgcaatga caaagtgctc agtagtggaa ggcaggctgt gaccagtctg cctgctcctt 2520 actataattg tgaggatttg ttactggaac agtacatgga ggcctgacct tgtgggggca 2580 cagggtggaa ccttagctga atatagtgtg tgtctcaaga ggaagtcagg gtactagctc 2640 agtgctcaat ctccaggtac tatatataca tttgcccgtt ttatctctaa tgtgaaataa 2700 ccactttett ggtttggtca ccccgatece eeggtettet getgtateta gaacagtgae 2820 tataaatgat gtatgggaat agtgtttcca tatgatctgt tgtctggagt atatgctaca 2880 tgttcattta ctgtacaaaa acccagtgca gctgatgatg caaagcagtc tctctctgtg 2940 tacagtgccc cacctattta aaaatcacgt acttgcccag aacactgtga aacacttaac 3000 ataagaacaa acgcagcgtc tggattcttt ccaaggagag cagctttctc cacaggaaca 3060 cagtaacaaa agaggtccgc cgccatccac acccagccaa gacacctcag aggccatagg 3120 gacaacetee ttgetggeea acaeetgetg gageagggge acaggteeca geaactgate 3180 ctcagtggat gggtctgcag ccaaagcctt aatgggctct cttttgaagg ggaaagaaag 3240 aatttcaagc ttatgatatc caatattatt atagttgatg agttagtaaa ttccaaaaaa 3300 aaaagatgat tttatatgta tgacataaaa aaaatctttg taaagtgcgc aagtgcaata 3360 atttaaagag gtcttatctt tgcatttata aattataaat attgtacatg tgtgtaattt 3420 ttcatgtatt catttgcagt ctttgtattt aaaaaaactt tactgttatg tttgtataat 3480 agaacattaa tcatttatta taactcagac aaggtgtaaa taaattcata attcaaacag 3540 ccagtatata tgcatatatg ggtgttacat tgcaaaaatc tctatctttg ttctattcac 3600 atgettaaag aagtaagaaa tettttgtgg atatgtaatt atacatataa agtatatata 3660 tatgtatgat acatgaaata tatttagaaa tgttcataat tttaatggat attctttggt 3720 aaaatttttt tttttttt ttattccaga gattaaagac actagatctt taaccttgaa 3840

```
gggcaggcaa gaggtcggca atgctgtcaa catagaagtc agggaccatt ttcttcttga 3900
acatgcagtc actttcctga ttgctcttca catcctcaag gctccggaat tccgqqqqtq 3960
tggtgggctt tgatctcagg actctggagg cagaagcagg cagatctctg tgaatatgag 4020
gccagcctgc actacacaga gctccagacc agtcatggct acatcatgaa accctgtctc 4080
aaaaagaaaa taaaaactgt tgtgtttcta ccatagtgtt aaactcagag tctgagtaat 4140
gtegggetga catgeteggg tgtttaacat acetteaget ttgaegagge getgaacagt 4200
caaagtctgg ccttggggag cggtggctgt gtttgtgctc aagtccaccg tgaaatcctg 4260
attgtgaatt tggacaaccg tgtccttctt cttggccttc catgcaacct ccaacttcat 4320
gttggtcatt ttgtcaaaac actgtgtgat gtttttatca atatactgcc attccacata 4380
tgtagagatg tagtctgcct ggctttcctt ttctttagcc aatcgaatgc tcttgatcat 4440
gccctcaatc tcatctctag cttttatcac gtctctgcta attcctgaaa cttgaatcga 4500
agttttcttc tggttcatct caatggtgat gttcagttcc ttctgaatct cattcagttt 4560
ctcgtactcc tccatgtcaa agtcactgac acactcatcg tcattggtgt aggaaagctg 4620
ctctttggta atcagttcct ttagccagga gattgttttg ttcacactgt ctacccctga 4680
accacatacc tggaaaactg tgtgctctat tttcttttcc aaaaccaggg tgttcttttt 4740
gggggaagct tgcttgggaa agccaagaaa ggctaaagag aaaatggaaa ttaatgtttc 4800
ttttactccc ttcaacatca aggttaggaa tatgtatttc ataaaagcta acaactcaca 4860
ggcaatctta gacatcactg actgcttggc aggcgactgc ttggggggag ctggagagcc 4920
ttctctttct ttcatgttgt cgtaaaaaaa ttgcagaata tggggctgga agataacaac 4980
tttaactctc ttcacagcct gcactgattt tttctggaca aattcttcaa tggcatctat 5040
tatcgctttt gctactacgt ttgggtcctg ttgagcattt ccttcaaaaa caaaaaagc 5100
acatttttaa aaagtcaagg ttaagatcca cctgcaaaaa aaagctgcaa tataagcgag 5160
gaattctagt tgtcacagga aataaaaatg tctgttccca ctataatcaa tgtagactga 5220
taatattatg ccagcaaata gttttgaagt cctaggcaca gtgggaggag gttttgttcc 5280
acgctgttca taagccaata ccccagcaaa agaccttaaa ggacaacttg taatttggga 5340
cattcacate tgteetette atetgatetg geteecagtg teacteteta acaeggteet 5400
tagagggaca atttateeet geetetgett gatettatge atgtatetgt attetteeag 5460
ccatccctgg cgacctgatt tttctaaggc acccaaaact gtaagctact tcttataatc 5520
tataattetg ageatattag ttageetgag cetecaggat atetttette cetataetea 5580
gtccagtttt agctgcccag aaggattcaa agctgatcta cgagtagatc actcctgtct 5640
acagettgtt ceagatettg ttteteaage eetggaagee ateageeagg taagattgta 5700
aaacaatccc tttctaatca tgggtgtggc ccaaagtgaa tggccggaat tc
                                                                  5752
```

<210> 3

<211> 475

<212> DNA

<213> mouse

<400> 3

tgtatgggaa tagtgtttcc atatgatctg ttgtctggag tatatgctac atgttcattt 60

```
actgtacaaa aacccagtgc agctgatgat gcaaagcagt ctctctctgt gtacagtgcc 120
ccacctattt aaaaatcacg tacttgccca gaacactgtg aaacacttaa cataagaaca 180
aacgcagcgt ctggattctt tccaaggaga gcagctttct ccacaggaac acagtaacaa 240
aagaggtccg ccgccatcca cacccagcca agacacctca gaggccatag ggacaacctc 300
cttgctggcc aacacctgct ggagcagggg cacaggtccc agcaactgat cctcagtgga 360
tgggtctgca gccaaagcct taatgggctc tcttttgaag gggaaagaaa gaatttcaag 420
cttatgatat ccaatattat tatagttgat gagttagtaa attccaaaaa aaaaa
<210> 4
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:primer
<400> 4
                                                                    20
agggctgtca atcatgctgg
<210> 5
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:primer
<400> 5
                                                                   20
aaactcacgg tcggtgcagc
<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:probe
```

WO 01/24781	PCT/CA00/01188
<400> 6	
attaaccctc actaaatgct gtat	24
<210> 7	
<211> 30	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:probe	•
<400> 7	
cattatgctg agtgatatct ttttttttcg	30
<210> 8	
<211> 38	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:probe	
<400> 8	
gaacatgtag catatactcc agacaacaga tcatatgg	38
<210> 9	
<211> 32	
<212> DNA	
<213> Artificial Sequence	•
<220>	
<223> Description of Artificial Sequence:probe	
<400> 9	
cagcttctcc acaggaacac agtaacaaag ag	32
<210> 10	
<211> 35	
<212> DNA	

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 10

ctatttcaca agagactgac cagccaataa atctc

35

<210> 11

<211> 7581

<212> DNA

<213> Unknown

<400> 11

cgcccgggca ggtctgttgg agggcagttg gtcaacctga ccagagagag ctgagctgga 60 agaccccact gatggtgtgc tgcctttcag tccaggaaga aagaaaggaa ggattctgag 120 gatttgggca aagccacatt cctggagaag tctgtatact gatgccaaac ccaagagctg 180 agctgctgat gaggcccagg gagtagccca cgcgccctga gctgttggct agcaaggcct 240 tcctgctcca tgtggcatgg aaaaattata tggtttgacg gatgaaaagg tgaaggccta 300 tetttetete catececagg tattagatga atttgtttet gaaagtgtta gtgeagagae 360 tgtggaaaag tggctgaaga ggaaaaccaa caaagcaaaa gatgaaccat ctcccaagga 420 agtcagcagg taccaggata cgaatatgca gggagtcgtg tacgagctga acagctacat 480 agagcagcgc ctggacacgg gcggggacaa ccacctgctc ctctatgagc tcagcagcat 540 catcaggata gccacaaaag ccgacggatt tgcactgtac ttccttggag agtgcaataa 600 tagcctgtgt gtgttcatac cacccgggat gaaggaaggc caaccccggc tcatccctgc 660 agggcccatc acccagggta ccaccatctc tgcctacgtg gccaagtcta ggaagacgtt 720 gttggtagag gatateettg gggatgageg attteetega ggtaetggee tggaateagg 780 aacccgcatc cagtctgttc tttgcttgcc cattgtcact gccattggag acttgattgg 840 catcettgaa etgtacagge actggggeaa agaggeette tgeeteagee atcaggaggt 900 tgcaacagec aatettgett gggetteegt ageaatacae caggtgcagg tgtgtagagg 960 tctcgccaaa cagaccgaac tgaatgactt cctactcgac gtatcaaaga catactttga 1020 taacatagtt gccatagact ctctacttga acacatcatg atatatgcaa aaaatctagt 1080 gaacgccgac cgctgcgcgc tettccaggt ggaccacaag aacaaggagc tgtactcgga 1140 cctgtttgac attggggagg agaaggaggg gaagcccatc ttcaagaaga ccaaggagat 1200 cagattttcc attgagaaag ggattgctgg tcaagtggca agaacaggcg aagtcttgaa 1260 cattcccgat gcctacgcgg accctcgctt taacagggag gtggacctgt acacaggcta 1320 caccacgagg aacattctgt gtatgcccat agtgagccga ggcagcgtga ttggcgtggt 1380 gcagatggtg aacaagatca gcggtagcgc cttctccaag acagacgaga acaacttcaa 1440 gatgtttgct gtcttctgcg cactggcctt gcactgtgct aacatgtacc acaggatccg 1500

ccactcag	aa tgcatctac	a gggttaccat	ggagaagctt	tcctaccaca	gcatctgcac	1560
ctccgagg	ag tggcaaggc	c tcatgcgctt	caacctacca	gcacgcatct	gccgggacat	1620
cgagctat	tc cactttgac	a ttggtccttt	cgagaacatg	tggcctggga	tctttgtcta	1680
catgatec	at cggtcttgt	g ggacatectg	ttttgaactt	gaaaaattgt	gccgttttat	1740
catgtctg	tg aagaagaac	t atcggcgggt	tccttaccac	aactggaa <u>g</u> c	atgcagtcac	1800
ggtggcac	ac tgcatgtat	g ccatacttca	aaacaacaat	ggcctcttca	cagacctcga	1860
gcgcaaag	gc ctgctaatt	g <mark>cgtgt</mark> ctgtg	ccatgacctg	gaccacaggg	gcttcagtaa	1920
cagctacc	tg cagaagtte	g accaccccct	ggcggcgctg	tactccacct	ccaccatgga	1980
gcaacacc	ac ttctcccag	a cggtgtccat	ccttcagctg	gaagggcaca	atatcttctc	2040
caccctgag	ge tecagegag	t acgagcaggt	gctggagatc	atccgcaaag	ccatcatcgc	2100
caccgacct	c gccctatact	t ttgggaacag	gaagcagttg	gaggagatgt	accagacagg	2160
gtcgctgaa	ac ctccacaac	c agtcccatcg	agaccgtgtċ	atcggcttga	tgatgactgc	2220
ctgtgatct	t tgctctgtg	a ccaaactatg	gccagttaca	aaattgacag	cgaatgatat	2280
atatgcaga	a ttctgggctg	g agggtgatga	gatgaagaag	ctgggcatac	agcccattcc	2340
tatgatgga	ac agagacaago	gagatgaagt	ccctcaaggg	cagctcggat	tctacaatgc	2400
tgtggccat	t ccctgctata	a ccaccttgac	gcagatcctc	ccacccacag	agcctctgct	2460
gaaggcctg	gc agggataaco	tcaatcagtg	ggagaaggta	attcgcgggg	aagagacagc	2520
aatgtggat	t tcaggcccag	geceggegee	tagcaagagc	acacctgaga	agctgaacgt	2580
gaaggttga	ia gactgatcct	gaagtgacgt	cctgatgtct	gcccagcaac	cgactcaacc	2640
tgcttctgt	g acttcgttct	tttgtttc	aaggggtgaa	aaccccctgt	cagaaggtac	2700
cgtcgcata	t ccatgtgaag	, cagacgactc	cctgcttgcc	gcacacacct	cggacagtga	2760
gcaacccag	g ctctgccgtg	ttcagacgtc	ggctactccg	tggctccacc	tgacctccga	2820
atgctattt	g ctcccaggcc	agcactgcac	tgtctggagg	gggcagagac	cacaggagag	2880
gttcttgcc	t gcatcctccc	atgagggtgt	ggccagttcc	ctagttctgt	gccatgctgc	2940
tgcttggtg	g cattggttag	gaatgggaca	cacgcccctt	gttgtgaagt	ttacatgtga	3000
ccttcttat	a ggttaactga	gtttgtggcc	tggacacatg	taatgaaggt	cacagtccac	3060
aggtgacag	a gaaatccaaa	ctgttgatta	caggtgcact	acaggtatgc	tctttcagtc	3120
tatctgggg	g cacataggtg	agtctgctcc	actcagaann	aagcatacct	ctgccctcat	3180
ccaggggac	a cagggtacat	cccaggcatc	ggggaactga	agctctcact	tcaaaccatg	3240
tcaaagaat	t aaaacacctc	ccctcccct	cactgtagcc	ttcgacaact	gcgccaatcc	3300
ctttataca	a agaaaataaa	agtaaggcat	ațaaatttcc	tccagcaagc	aaatcttgtg	3360
ggtaaaaaa	a aagcatgtga	atnntaacaa	cntctanant	ntcncngnat	gttatggcag	3420
aattttagt	c acgtccaaaa	caaaaagatt	attccagaag	atacctcatc	ctatgcctga	3480
aaggctcca	c agcatggcgt	ccgtctccca	gggttctgat	ccgtctcctc	acggtgcaat	3540
caggcagga	c agagaggagg	gctgcagggc	taccacattg	acccagaagg	tatctcctct	3600
	g acatccataa					
	a gccaggacac					
	g gattttcagg		_			
gagccttga	g aaggagagac	tgaccagaaa	cactcactca	gcactctgca	ggagcaggag	3840

aagatacttt aagatgaatc ttggatagat tttgatacac ccaataccat acacacagga 3900 gcttggcatt tgcaaagtct attcagtttc cttccgcgct ctgacccacg gttgtagcgg 3960 agtgggctga acactgtaac actgtacatg cgatttcccc atgggcttct aaaatgtcac 4020 catctcctcc cctgctgtgt cctactccat ttactggtta caaggtgatg tcaacaagag 4080 aagctatcac aacaccaggg ctgtgcacac gtgcacacac atgtatgcac aagcacacag 4140 atgtatgtac agcacacaca cacacaca ccccaaaagg agagaaaagg aagaaaacat 4200 ttataaaaag cgacagctac cccatatcaa aatagtcttt cctgtaggaa acaggagctc 4260 tccataagga attatcatga gtgtgttctc ccatcagtgc actctcccag gggtgctcac 4320 tgaagetggt ccaertetat aaacaggtga caetggetge agcaaaaage cattegatee 4380 acacaaattg atcttctatc atcttggaat ctgaattgca gggaggagca gyatgtaaga 4440 cgaccgttta attcaggcat tccgaaggca tgagcgcatg gattctrtca ccaagcgtat 4500 aaaaggaccc tggcattggg aaacctatga cggactgttt ttgctgtaga agtagggatt 4560 ttacagaagt ctccttgrat ttgccctgcc tggggcagtt ttgcagagga acctgccaga 4620 gatttattgg ctggtcagtc tcttgtgaaa tagtatcatg tgagaaacag tttgtagaaa 4680 aaaactatac ctgggaagac ctttgcaaca ttgttccttc catgggccaa gactcagtta 4740 ggaggcataa atctgcccgg aataaactag gccaggatac agccatgttt agttaataat 4800 ttggttttag aattcacaca ggcaggattg gtttttttgt gtcttggcaa gtggagcata 4860 tttaacatac aggcatggga atcctgcctc ttagcttttc ccaccctctt gtctcaccaa 4920 gttttttctc tccaaaggtt tccaggaatt tctcattaat ggctgatgca aacttagtga 4980 ataataatga atataaacaa tgctcacctc accaaaatta tattatttgc agtcatttgt 5040 gataacacaa attttatcgc aatggttatt atttaatttg tggccacaca ctgtggttat 5100 cttttgttgt ggttgtttct gagaaaatgt tcttggatat gtaagtgcca ataccagtgt 5160 gaagtattga tcccgggcag caaaatacag cctaaggttt gtaaacatca attctatctc 5220 agttcatcag agggcctgag aagctgcggg gcagtgtaaa gtaaagtatg ctgggctggt 5280 ggtggtcagc ctccccttgc caagaagaga gcaattgaat cctgtcccca gctccctcca 5340 cgcctgaaga gtgaccagtg ctggcccgac ggatcgctga gatattctcc cataatggca 5400 aaaaaatagg cagtttgatg tgacctgttt agtgtggctc tcctcttttg agcatgtgtt 5460 agcattttta ttttatactc atccagtgaa ctctgctctt ccaagtgtgt tcatgtatgt 5520 gctagatata ttagcacage etgeettetg etgeacaaeg cettagagae eeggeettte 5580 aatgagetta gettgtgete tgtttetget etettaggte taaactatgg tgteagtttt 5640 aatagaacaa aagtatgcat cttgccttgg cttgagcctt ttcgttttca atgctgactt 5700 ctcccctttc tctcctgtgc tcaccttacc tttccagagt gtaagggaca acttttaagg 5760 aggegtgtee etggtagggg catecetgtt caccaggtge etgteateae eccaettgae 5820 tgacatctac cctggtgact atgggttcct cttgtttgta gggaacggtg gctccaggtg 5880 gaggcatcaa tetgttgggt tetggtteee ggetgeettt ggttttgaaa gtetettete 5940 tgtatattcc taccetgeat ttgetttgtg tggtgetgat getgtggeag taggatettg 6000 gatgactete cateagteae agacteecee tgttgeaaag tgteaggetg actegaeagt 6060 caccgtaaaa tetgagteag teacacaeag getgteagee aeggetteea ettgeatgge 6120 tattctattt tcacacgtga gtttctgttg ctggctggct gactggcatt atctatgcta 6180

agttgaaatc aggagtgtgc ccagcagagc ccatcattct cactgtcttt gaaacaaagc 6240 tgtacggttt gatcgatgaa cgtatttaaa gcatttcatg caatgacaaa gtgctcagta 6300 gtggaaggca ggctgtgacc agtctgcctg ctccttacta taattgtgag gatttgttac 6360 tggaacagta catggaggcc tgaccttgtg ggggcacagg gtggaacctt agctgaatat 6420 agtgtgtgtc tcaagaggaa gtcagggtac tagctcagtg ctcaatctcc aggtactata 6480 tatacatttg cccgttttat ctctaatgtg aaataaatcc ccaaacactt gtttatcgtg 6540 tagcgtacct aaaagactat tctattatgg gtgtccccac tttcttggtt tggtcacccc 6600 gateceeegg tettetgetg tatetagaae agtgaetata aatgatgtat gggaatagtg 6660 tttccatatg atctgttgtc tggagtatat gctacatgtt catttactgt acaaaaaccc 6720 agtgcagctg atgatgcaaa gcagtctctc tctgtgtaca gtgccccacc tatttaaaaa 6780 tcacgtacaa ncccagaaca ctgtgaaaca cttaacataa gaaacaaacg cagcgtctgg 6840 attettteea aggagageag ettteteeae aggaacaeag taacaaaaga ggteegeege 6900 catccacacc cagccaagac acctcagagg ccatagggac aacctccttg ctggccaaca 6960 cctgctggag cagggcacag gtcccagcaa ctgatcctca gtggatgggt ccgcagtcaa 7020 agccttaatg ggctctcttt tgaaggggaa agaaannttt caagcttatg atatccaaca 7080 ttattatagt tgatgagtta gtaaattccg aaaaaaaaag atgattttat atgtatgaca 7140 taaaaaaaat ctttgtaaag tgcgcaagtg caataattta aagaggtctt atctttgcat 7200 ttataaatta taaatattgt acatgtgtgt aatttttcat gtattcattt gcagtctttg 7260 tatttaaaaa aactttactg ttatgtttgt ataatagaac attaatcatt tattataact 7320 cagacaaggt gtaaataaat tcataattca aacagccagt atatatgcat atatgggtgt 7380 tacattgcaa aaatctctat ctttgttcta ttcacatgct taaagaagta agaaatcttt 7440 tgtggatatg taattataca tataaagtat atatatatgt atgatacatg aaatatattt 7500 agaaatgttc ataattttaa tggatattct ttggtgtgaa taattgaata caacattttt 7560 7581 aaaatgaaaa aaaaaaaaaa c

<210> 12

<211> 7618

<212> DNA

<213> mouse

<400> 12

egeceggea ggtetgttgg agggeagttg gteaacetga ceagagaga etgagetgga 60 agaceceaet gatggtgte tgeetteag teeaggaaga aagaaaggaa ggattetgag 120 gatttgggea aagecaeatt eetggagaag tetgtataet gatgeeaaae eeaaggeetg 180 agetgetgat gaggeeeagg gagtageeea egegeeetga getgttgget ageaaggeet 240 teetgeteea tgtggeatgg aaaaattata tggtttgaeg gatgaaaagg tgaaggeeta 300 tettteetee eateeeagg tattagatga atttgttet gaaagtgtta gtgeagagae 360 tgtgggaaaag tggetgaaga ggaaaaceaa caaageaaaa gatgaaeeat eteeeaagga 420 agteageagg taceaggata egaatatgea gggagtegtg taegagetga acagetaeat 480

agagcagege ctggacaegg geggggacaa ceacetgete etetatgage teagcageat 540 catcaggata gccacaaaag ccgacggatt tgcactgtac ttccttggag agtgcaataa 600 tagcctgtgt gtgttcatac cacccgggat gaaggaaggc caaccccggc tcatccctgc 660 agggcccatc acccagggta ccaccatctc tgcctacgtg gccaagtcta ggaagacgtt 720 gttggtagag gatateettg gggatgageg attteetega ggtaetggee tggaateagg 780 aacccgcatc cagtctgttc tttgcttgcc cattgtcact gccattggag acttgattgg 840 catcettgaa etgtacagge aetggggeaa agaggeette tgeeteagee atcaggaggt 900 tgcaacagcc aatcttgctt gggcttccgt agcaatacac caggtgcagg tgtgtagagg 960 tctcgccaaa cagaccgaac tgaatgactt cctactcgac gtatcaaaga catactttga 1020 taacatagtt gccatagact ctctacttga acacatcatg atatatgcaa aaaatctagt 1080 gaacgccgac cgctgcgcgc tcttccaggt ggaccacaag aacaaggagc tgtactcgga 1140 cctgtttgac attggggagg agaaggaggg gaagcccatc ttcaagaaga ccaaggagat 1200 cagattttcc attgagaaag ggattgctgg tcaagtggca agaacaggcg aagtcttgaa 1260 cattecegat geetaegegg accetegett taacagggag gtggacetgt acacaggeta 1320 caccacgagg aacattetgt gtatgeecat agtgageega ggeagegtga ttggegtggt 1380 gcagatggtg aacaagatca gcggtagcgc cttctccaag acagacgaga acaacttcaa 1440 gatgtttgct gtcttctgcg cactggcctt gcactgtgct aacatgtacc acaggatccg 1500 ccactcagaa tgcatctaca gggttaccat ggagaagctt tcctaccaca gcatctgcac 1560 ctccgaggag tggcaaggcc tcatgcgctt caacctacca gcacgcatct gccgggacat 1620 cgagctattc cactttgaca ttggtccttt cgagaacatg tggcctggga tctttgtcta 1680 catgatccat cggtcttgtg ggacatcctg ttttgaactt gaaaaattgt gccgttttat 1740 catgtctgtg aagaagaact atcggcgggt tccttaccac aactggaagc atgcagtcac 1800 ggtggcacac tgcatgtatg ccatacttca aaacaacaat ggcctcttca cagacctcga 1860 gcgcaaaggc ctgctaattg cgtgtctgtg ccatgacctg gaccacaggg gcttcagtaa 1920 cagetacetg cagaagtteg accaeeeet ggeggegetg taeteeaeet ecaeeatgga 1980 gcaacaccac ttctcccaga cggtgtccat ccttcagctg gaagggcaca atatcttctc 2040 caccetgage tecagegagt aegageaggt getggagate ateegeaaag ceateatege 2100 caccgacctc gccctatact ttgggaacag gaagcagttg gaggagatgt accagacagg 2160 gtcgctgaac ctccacaacc agtcccatcg agaccgtgtc atcggcttga tgatgactgc 2220 ctgtgatctt tgctctgtga ccaaactatg gccagttaca aaattgacag cgaatgatat 2280 atatgcagaa ttctgggctg agggtgatga gatgaagaag ctgggcatac agcccattcc 2340 tatgatggac agagacaagc gagatgaagt ccctcaaggg cagctcggat tctacaatgc 2400 tgtggccatt ccctgctata ccaccttgac gcagatectc ccacccacag agectctgct 2460 gaaggeetge agggataace teaateagtg ggagaaggta attegegggg aagagacage 2520 aatgtggatt teaggeceag geeeggegee tageaagage acaeetgaga agetgaaegt 2580 gaaggttgaa gactgateet gaagtgaegt eetgatgtet geeeageaac egacteaace 2640 tgcttctgtg acttcgttct ttttgttttc aaggggtgaa aaccccctgt cagaaggtac 2700 cgtcgcatat ccatgtgaag cagacgactc cctgcttgcc gcacacacct cggacagtga 2760 gcaacccagg ctctgccgtg ttcagacgtc ggctactccg tggctccacc tgacctccga 2820

atgctatttg ctcccaggcc agcactgcac tgtctggagg gggcagagac cacaggagag 2880 gttcttgcct gcatcctccc atgagggtgt ggccagttcc ctagttctgt gccatgctgc 2940 tgcttggtgg cattggttag gaatgggaca cacgcccctt gttgtgaagt ttacatgtga 3000 cettettata ggttaactga gtttgtggcc tgggacacat gtaatgaagg tcacagtcca 3060 caggtgacag agaaatccaa actgttgatt acaggtgcac tacaggtatg ctctttcagt 3120 ctatctgggg gcacataggt gagtctgctc cactcagaag gaagcatacc tctsccctca 3180 tccaggggac acagggtaca tcccaggcat cggggaactg aagctctcac ttcaaaccat 3240 gtcaaagaat taaaacacct cccctccccc tcactgtagc cttcggcaac tgcgccaatc 3300 cctttataca aagaaaatat aagtaaggca tataaatttc ctccagcaag caaatcttgt 3360 gggtaaaaaa aaaaaatgtg aattttaaca acctctatat tttcactgta tgttatggca 3420 gaattttagt cacgtccaaa acaaaagatt attccagaag atacctcatc ctatgcctga 3480 aagctccaca gcatggcgtc cgtctcccag ggttctgatc cgtctcctca cggtgcaatc 3540 aggcaggaca ggaggaggtg cagggctacc acattgaccc agatggtatc tcctctcacc 3600 attcagacat ccataaggaa tgccaaatgc tgtattgaat agttctcctg tgtgactttc 3660 tagagaagee aggacaceee tgageettte etgggaacte etaaggaagt cacaggttea 3720 caccgtgggg attttcagga tagcatggag accagagaat cccggttcgg ttgttctcac 3780 teggtgagee ttgagaagga agagaetgae cagaaacaet caeteageae tetggeagga 3840 gcaggagaag atactttaag atgaatcttt gggatagatt ttgatacacc caataccata 3900 cacacaggag cttggcattt gcaaagtcta ttcagtttcc ttccacactc tgacccacgg 3960 ttgtagcgga gtgggctgaa cactgtaaca ctgtacatgc gatttcccca tgggcttcta 4020 aaatgtcacc atctcctccc ctgctgtgtc ctactccatt tactggttac aaggtgatgt 4080 caacaagaga agctatcaca acaccagggc tgtgcacacg tgcacacaca tgtatgcaca 4140 agcacacaga tgtatgtaca gcacacacac acacacacac cccaaaagga gagaaaagga 4200 agaaaacatt tataaaaagc gacagctacc cccatattca aaaatagttc ttttccctgt 4260 agggaaacag gtagctctcc ataaggaaat tatcatgagt gtgttctccc atcagtgcac 4320 ttctcccagg ggtgctcact gaagctggtc cacgtctata aacaggtgac actggctgca 4380 gcaaaaagcc attcgatcca cacaaattga tettetatca tettggaate tgaattgcag 4440 ggaggagcag catgtaagac gaccgtttaa ttcaggcatt ccgaaggcat gagcgcatgg 4500 attctgtcac caagcgtata aaaggaccct ggcattggga aacctatgac ggactgtttt 4560 tgctgtagaa gtagggattt tacagaagtc tccttggatt tgccctgcct ggggcagttt 4620 tgcagaggaa cctgccagag atttattggc tggtcagtct cttgtgaaat agtatcatgt 4680 gagaaacagt ttgtagaaaa aaactatacc tgggaagacc tttgcaacat tgttccttcc 4740 atgggccaag actcagttag gaggcataaa tctgcccgga ataaactagg ccaggataca 4800 gccatgttta gttaataatt tggttttaga attcacacag gcaggattgg ttttttttgtg 4860 tettggeaag tggageatat ttaacataea ggeatgggaa teetgeetet tagettttee 4920 caccetettg teteaceaag tittitetet ceaaaggitti eeaggaatit eteattaatg 4980 gctgatgcaa acttagtgaa taataatgaa tataaacaat gctcacctca ccaaaattat 5040 attatttgca gtcatttgtg ataacacaaa ttttatcgca atggttatta tttaatttgt 5100 ggccacacac tgtggttatc ttttgttgtg gttgtttctg agaaaatgtt cttggatatg 5160

taagtgccaa taccagtgtg aagtattgat cccgggcagc aaaatacagc ctaaggtttg 5220 taaacatcaa ttctatctca gttcatcaga gggcctgaga agctgcgggg cagtgtaaag 5280 taaagtatgc tgggctggtg gtggtcagcc teceettgec aagaagagag caattgaatc 5340 ctgtccccag ctccctccac gcctgaagag tgaccagtgc tggcccgacg gatcgctgag 5400 atattetece ataatggeaa aaaaatagge agtttgatgt gacetgttta gtgtggetet 5460 cctcttttga gcatgtgtta gcatttttat tttatactca tccagtgaac tctgctcttc 5520 caagtgtgtt catgtatgtg ctagatatat tagcacagec tgeettetge tgeacaaege 5580 cttagagacc cggcctttca atgagcttag cttgtgctct gtttctgctc tcttaggtct 5640 aaactatggt gtcagtttta atagaacaaa agtatgcatc ttgccttggc ttgagccttt 5700 tegttttcaa tgctgacttc teceetttet eteetgtget eacettacet ttecagagtg 5760 taagggacaa cttttaagga ggcgtgtccc tggtaggggc atccctgttc accaggtgcc 5820 tgtcatcacc ccacttgact gacatctacc ctggtgacta tgggttcctc ttgtttgtag 5880 ggaacggtgg ctccaggtgg aggcatcaat ctgttgggtt ctggttcccg gctgcctttg 5940 gttttgaaag tetettetet gtatatteet accetgeatt tgetttgtgt ggtgetgatg 6000 ctgtggcagt aggatcttgg atgactetec atcagteaca gacteceet gttgcaaagt 6060 gtcaggctga ctcgacagtc accgtaaaat ctgagtcagt cacacacagg ctgtcagcca 6120 eggetteeae ttgeatgget attetatttt cacaegtgag tttetgttge tggetggetg 6180 actggcatta tctatgctaa gttgaaatca ggagtgtgcc cagcagagcc catcattctc 6240 actgtctttg aaacaaagct gtacggtttg atcgatgaac gtatttaaag catttcatgc 6300 aatgacaaag tgctcagtag tggaaggcag gctgtgacca gtctgcctgc tccttactat 6360 aattgtgagg atttgttact ggaacagtac atggaggcct gaccttgtgg gggcacaggg 6420 tggaacetta getgaatata gtgtgtgtet caagaggaag teagggtaet ageteagtge 6480 tcaatctcca ggtactatat atacatttgc ccgttttatc tctaatgtga aataaatccc 6540 caaacacttg tttatcgtgt agcgtaccta aaagactatt ctattatggg tgtccccact 6600 ttettggttt ggteaceeeg ateceeeggt ettetgetgt atetagaaca gtgaetataa 6660 atgatgtatg ggaatagtgt ttccatatga tctgttgtct ggagtatatg ctacatgttc 6720 atttactgta caaaaaccca gtgcagctga tgatgcaaag cagtctctct ctgtgtacag 6780 tgccccacct atttaaaaat cacgtacaan cccagaacac tgtgaaacac ttaacataag 6840 aaacaaacgc agcgtctgga ttctttccaa ggagagcagc tttctccaca ggaacacagt 6900 aacaaaagag gtccgccgcc atccacaccc agccaagaca cctcagaggc catagggaca 6960 accteettge tggeeaacae etgetggage agggeacagg teccageaae tgateeteag 7020 tggatgggtc cgcagtcaaa gccttaatgg gctctctttt gaaggggaaa gaaanntttc 7080 aagettatga tateeaacat tattatagtt gatgagttag taaatteega aaaaaaaaga 7140 tgattttata tgtatgacat aaaaaaaatc tttgtaaagt gcgcaagtgc aataatttaa 7200 agaggtetta tetttgeatt tataaattat aaatattgta catgtgtgta attttteatg 7260 tattcatttg cagtctttgt atttaaaaaa actttactgt tatgtttgta taatagaaca 7320 ttaatcattt attataactc agacaaggtg taaataaatt cataattcaa acagccagta 7380 tatatgcata tatgggtgtt acattgcaaa aatctctatc tttgttctat tcacatgctt 7440 aaagaagtaa gaaatctttt gtggatatgt aattatacat ataaagtata tatatatgta 7500

tgatacatga aatatatta gaaatgttca taattttaat ggatattctt tggtgtgaat 7560 aattgaatac aacattttta aaatgaaaaa aaaaaaaaa aaaaaaaaa 7618